

## Answers

## CHAPTER 1

## Prior knowledge check

- 1 a  $2m^2n + 3mn^2$       b  $6x^2 - 12x - 10$   
 2 a  $2^8$       b  $2^4$       c  $2^6$   
 3 a  $3x + 12$       b  $10 - 15x$       c  $12x - 30y$   
 4 a 8      b  $2x$       c  $xy$   
 5 a  $2x$       b  $10x$       c  $\frac{5x}{3}$

## Exercise 1A

- 1 a  $x^7$       b  $6x^5$       c  $k$       d  $2p^2$   
 e  $x$       f  $y^{10}$       g  $5x^2$       h  $p^2$   
 i  $2a^3$       j  $2p$       k  $6a^9$       l  $3a^2b^3$   
 m  $27x^8$       n  $24x^{11}$       o  $63a^{12}$       p  $32y^6$   
 q  $4a^6$       r  $6a^{12}$
- 2 a  $9x - 18$       b  $x^2 + 9x$   
 c  $-12y + 9y^2$       d  $xy + 5x$   
 e  $-3x^2 - 5x$       f  $-20x^2 - 5x$   
 g  $4x^2 + 5x$       h  $-15y + 6y^3$   
 i  $-10x^2 + 8x$       j  $3x^3 - 5x^2$   
 k  $4x - 1$       l  $2x - 4$   
 m  $9d^2 - 2c$       n  $13 - r^2$   
 o  $3x^3 - 2x^2 + 5x$       p  $14y^2 - 35y^3 + 21y^4$   
 q  $-10y^2 + 14y^3 - 6y^4$       r  $4x + 10$   
 s  $11x - 6$       t  $7x^2 - 3x + 7$   
 u  $-2x^2 + 26x$       v  $-9x^3 + 23x^2$
- 3 a  $3x^3 + 5x^5$       b  $3x^4 - x^6$       c  $\frac{x^3}{2} - x$   
 d  $4x^2 + \frac{5}{2}$       e  $\frac{7x^6}{5} + x$       f  $3x^4 - \frac{5x^2}{3}$

## Exercise 1B

- 1 a  $x^2 + 11x + 28$   
 b  $x^2 - x - 6$   
 c  $x^2 - 4x + 4$   
 d  $2x^2 + 3x - 2xy - 3y$   
 e  $4x^2 + 11xy - 3y^2$   
 f  $6x^2 - 10xy - 4y^2$   
 g  $2x^2 - 11x + 12$   
 h  $9x^2 + 12xy + 4y^2$   
 i  $4x^2 + 6x + 16xy + 24y$   
 j  $2x^2 + 3xy + 5x + 15y - 25$   
 k  $3x^2 - 4xy - 8x + 4y + 5$   
 l  $2x^2 + 5x - 7xy - 4y^2 - 20y$   
 m  $x^2 + 2x + 2xy + 6y - 3$   
 n  $2x^2 + 15x + 2xy + 12y + 18$   
 o  $13y - 4x + 12 - 4y^2 + xy$   
 p  $12xy - 4y^2 + 3y + 15x + 10$   
 q  $5xy - 20y - 2x^2 + 11x - 12$   
 r  $22y - 4y^2 - 5x + xy - 10$
- 2 a  $5x^2 - 15x - 20$   
 b  $14x^2 + 7x - 70$   
 c  $3x^2 - 18x + 27$   
 d  $x^3 - xy^2$   
 e  $6x^3 + 8x^2 + 3x^2y + 4xy$   
 f  $x^2y - 4xy - 5y$   
 g  $12x^2y + 6xy - 8xy^2 - 4y^2$   
 h  $19xy - 35y - 2x^2y$   
 i  $10x^3 - 4x^2 + 5x^2y - 2xy$   
 j  $x^3 + 3x^2y - 2x^2 + 6xy - 8x$

- k  $2x^2y + 9xy + xy^2 + 5y^2 - 5y$   
 l  $6x^2y + 4xy^2 + 2y^2 - 3xy - 3y$   
 m  $2x^3 + 2x^2y - 7x^2 + 3xy - 15x$   
 n  $24x^3 - 6x^2y - 26x^2 + 2xy + 6x$   
 o  $6x^3 + 15x^2 - 3x^2y - 18xy^2 - 30xy$   
 p  $x^3 + 6x^2 + 11x + 6$   
 q  $x^3 + x^2 - 14x - 24$   
 r  $x^3 - 3x^2 - 13x + 15$   
 s  $x^3 - 12x^2 + 47x - 60$   
 t  $2x^3 - x^2 - 5x - 2$   
 u  $6x^3 + 19x^2 + 11x - 6$   
 v  $18x^3 - 15x^2 - 4x + 4$   
 w  $x^3 - xy^2 - x^2 + y^2$   
 x  $8x^3 - 36x^2y + 54xy^2 - 27y^3$
- 3  $2x^2 - xy + 29x - 7y + 24$   
 4  $4x^3 + 12x^2 + 5x - 6 \text{ cm}^3$   
 5  $a = 12, b = 32, c = 3, d = -5$

## Challenge

$$x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

## Exercise 1C

- 1 a  $4(x + 2)$       b  $6(x - 4)$   
 c  $5(4x + 3)$       d  $2(x^2 + 2)$   
 e  $4(x^2 + 5)$       f  $6x(x - 3)$   
 g  $x(x - 7)$       h  $2x(x + 2)$   
 i  $x(3x - 1)$       j  $2x(3x - 1)$   
 k  $5y(2y - 1)$       l  $7x(5x - 4)$   
 m  $x(x + 2)$       n  $y(3y + 2)$   
 o  $4x(x + 3)$       p  $5y(y - 4)$   
 q  $3xy(3y + 4x)$       r  $2ab(3 - b)$   
 s  $5x(x - 5y)$       t  $4xy(3x + 2y)$   
 u  $5y(3 - 4z^2)$       v  $6(2x^2 - 5)$   
 w  $xy(y - x)$       x  $4y(3y - x)$
- 2 a  $x(x + 4)$       b  $2x(x + 3)$   
 c  $(x + 8)(x + 3)$       d  $(x + 6)(x + 2)$   
 e  $(x + 8)(x - 5)$       f  $(x - 6)(x - 2)$   
 g  $(x + 2)(x + 3)$       h  $(x - 6)(x + 4)$   
 i  $(x - 5)(x + 2)$       j  $(x + 5)(x - 4)$   
 k  $(2x + 1)(x + 2)$       l  $(3x - 2)(x + 4)$   
 m  $(5x - 1)(x - 3)$       n  $2(3x + 2)(x - 2)$   
 o  $(2x - 3)(x + 5)$       p  $2(x^2 + 3)(x^2 + 4)$   
 q  $(x + 2)(x - 2)$       r  $(x + 7)(x - 7)$   
 s  $(2x + 5)(2x - 5)$       t  $(3x + 5y)(3x - 5y)$   
 u  $4(3x + 1)(3x - 1)$       v  $2(x + 5)(x - 5)$   
 w  $2(3x - 2)(x - 1)$       x  $3(5x - 1)(x + 3)$
- 3 a  $x(x^2 + 2)$       b  $x(x^2 - x + 1)$   
 c  $x(x^2 - 5)$       d  $x(x + 3)(x - 3)$   
 e  $x(x - 4)(x + 3)$       f  $x(x + 5)(x + 6)$   
 g  $x(x - 1)(x - 6)$       h  $x(x + 8)(x - 8)$   
 i  $x(2x + 1)(x - 3)$       j  $x(2x + 3)(x + 5)$   
 k  $x(x + 2)(x - 2)$       l  $3x(x + 4)(x + 5)$
- 4  $(x^2 + y^2)(x + y)(x - y)$   
 5  $x(3x + 5)(2x - 1)$

## Challenge

$$(x - 1)(x + 1)(2x + 3)(2x - 3)$$

## Exercise 1D

- 1 a  $x^5$       b  $x^{-2}$       c  $x^4$       d  $x^3$   
 e  $x^5$       f  $12x^0 = 12$       g  $3x^{\frac{1}{2}}$       h 5x  
 i  $6x^{-1}$       j  $x^{\frac{5}{6}}$       k  $x^{\frac{17}{6}}$       l  $x^{\frac{1}{6}}$

- 2 a 5      b 729      c 3      d  $\frac{1}{16}$   
 e  $\frac{1}{3}$       f  $\frac{-1}{125}$       g 1      h 216  
 i  $\frac{125}{64}$       j  $\frac{9}{4}$       k  $\frac{5}{6}$       l  $\frac{64}{49}$
- 3 a  $8x^5$       b  $\frac{5}{x^2} - \frac{2}{x^3}$       c  $5x^4$   
 d  $\frac{1}{x^2} + 4$       e  $\frac{2}{x^3} + \frac{1}{x^2}$       f  $\frac{8}{27}x^6$   
 g  $\frac{3}{x} - 5x^2$       h  $\frac{1}{3x^2} + \frac{1}{5x}$
- 4 a 3      b  $\frac{16}{\sqrt[3]{x}}$
- 5 a  $\frac{x}{2}$       b  $\frac{32}{x^6}$

**Exercise 1E**

- 1 a  $2\sqrt{7}$       b  $6\sqrt{2}$       c  $5\sqrt{2}$       d  $4\sqrt{2}$   
 e  $3\sqrt{10}$       f  $\sqrt{3}$       g  $\sqrt{3}$       h  $6\sqrt{5}$   
 i  $7\sqrt{2}$       j  $12\sqrt{7}$       k  $-3\sqrt{7}$       l  $9\sqrt{5}$   
 m  $23\sqrt{5}$       n 2      o  $19\sqrt{3}$
- 2 a  $2\sqrt{3} + 3$       b  $3\sqrt{5} - \sqrt{15}$   
 c  $4\sqrt{2} - \sqrt{10}$       d  $6 + 2\sqrt{5} - 3\sqrt{2} - \sqrt{10}$   
 e  $6 - 2\sqrt{7} - 3\sqrt{3} + \sqrt{21}$       f  $13 + 6\sqrt{5}$   
 g  $8 - 6\sqrt{3}$       h  $5 - 2\sqrt{3}$   
 i  $3 + 5\sqrt{11}$
- 3  $3\sqrt{3}$

**Exercise 1F**

- 1 a  $\frac{\sqrt{5}}{5}$       b  $\frac{\sqrt{11}}{11}$       c  $\frac{\sqrt{2}}{2}$   
 d  $\frac{\sqrt{5}}{5}$       e  $\frac{1}{2}$       f  $\frac{1}{4}$   
 g  $\frac{\sqrt{13}}{13}$       h  $\frac{1}{3}$
- 2 a  $\frac{1 - \sqrt{3}}{-2}$       b  $\sqrt{5} - 2$       c  $\frac{3 + \sqrt{7}}{2}$   
 d  $3 + \sqrt{5}$       e  $\frac{\sqrt{5} + \sqrt{3}}{2}$       f  $\frac{(3 - \sqrt{2})(4 + \sqrt{5})}{11}$   
 g  $5(\sqrt{5} - 2)$       h  $5(4 + \sqrt{14})$       i  $\frac{11(3 - \sqrt{11})}{-2}$   
 j  $\frac{5 - \sqrt{21}}{-2}$       k  $\frac{14 - \sqrt{187}}{3}$       l  $\frac{35 + \sqrt{1189}}{6}$   
 m -1
- 3 a  $\frac{11 + 6\sqrt{2}}{49}$       b  $9 - 4\sqrt{5}$       c  $\frac{44 + 24\sqrt{2}}{49}$   
 d  $\frac{81 - 30\sqrt{2}}{529}$       e  $\frac{13 + 2\sqrt{2}}{161}$       f  $\frac{7 - 3\sqrt{3}}{11}$
- 4  $-\frac{7}{4} + \frac{\sqrt{5}}{4}$

**Mixed exercise**

- 1 a  $y^8$       b  $6x^7$       c  $32x$       d  $12b^9$   
 2 a  $x^2 - 2x - 15$       b  $6x^2 - 19x - 7$   
 c  $6x^2 - 2xy + 19x - 5y + 10$

- 3 a  $x^3 + 3x^2 - 4x$       b  $x^3 + 6x^2 - 13x - 42$   
 c  $6x^3 - 5x^2 - 17x + 6$
- 4 a  $15y + 12$       b  $15x^2 - 25x^3 + 10x^4$   
 c  $16x^2 + 13x$       d  $9x^3 - 3x^2 + 4x$   
 5 a  $x(3x + 4)$       b  $2y(2y + 5)$   
 c  $x(x + y + y^2)$       d  $2xy(4y + 5x)$   
 6 a  $(x + 1)(x + 2)$       b  $3x(x + 2)$   
 c  $(x - 7)(x + 5)$       d  $(2x - 3)(x + 1)$   
 e  $(5x + 2)(x - 3)$       f  $(1 - x)(6 + x)$   
 7 a  $2x(x^2 + 3)$       b  $x(x + 6)(x - 6)$   
 c  $x(2x - 3)(x + 5)$
- 8 a  $3x^6$       b 2      c  $6x^2$       d  $\frac{1}{2}x^{-3}$
- 9 a  $\frac{4}{9}$       b  $\pm \frac{3375}{4913}$
- 10 a  $\frac{\sqrt{7}}{7}$       b  $4\sqrt{5}$
- 11 a 21877  
 b  $(5x + 6)(7x - 8)$   
 When  $x = 25$ ,  $5x + 6 = 131$  and  $7x - 8 = 167$ ; both 131 and 167 are prime numbers.
- 12 a  $3\sqrt{2} + \sqrt{10}$       b  $10 + 2\sqrt{3} - 5\sqrt{5} - \sqrt{15}$   
 c  $24 - 6\sqrt{7} - 4\sqrt{2} + \sqrt{14}$
- 13 a  $\frac{\sqrt{3}}{3}$       b  $\sqrt{2} + 1$       c  $-3\sqrt{3} - 6$   
 d  $\frac{30 - \sqrt{851}}{-7}$       e  $7 - 4\sqrt{3}$       f  $\frac{23 + 8\sqrt{7}}{81}$
- 14 a  $b = -4$  and  $c = -5$       b  $(x + 3)(x - 5)(x + 1)$
- 15 a  $\frac{1}{4}x$       b  $256x^{-3}$
- 16  $\frac{5}{\sqrt{75} - \sqrt{50}} = \frac{1}{\sqrt{3} - \sqrt{2}} = \sqrt{3} + \sqrt{2}$
- 17  $-36 + 10\sqrt{11}$
- 18  $x(1 + 8x)(1 - 8x)$
- 19  $y = 6x + 3$
- 20  $4\sqrt{3}$
- 21  $3 - \sqrt{3}$  cm
- 22  $\frac{4 - 4x^{\frac{1}{2}} + x^{\frac{1}{4}}}{x^{\frac{1}{2}}} = 4x^{-\frac{1}{2}} - 4 + x^{\frac{1}{4}}$
- 23  $\frac{11}{2}$
- 24  $4x^{\frac{3}{2}} + x^2$ ,  $a = \frac{5}{2}$ ,  $b = 2$



**Challenge**



- a  $a - b$   
 b  $\frac{(\sqrt{1} - \sqrt{2}) + (\sqrt{2} - \sqrt{3}) + \dots + (\sqrt{24} - \sqrt{25})}{-1} = \sqrt{25} - \sqrt{1} = 4$

**CHAPTER 2**

**Prior knowledge check**

- 1 a  $x = -5$       b  $x = 3$   
 c  $x = 5$  or  $x = -5$       d 16 or 0  
 2 a  $(x + 3)(x + 5)$       b  $(x + 5)(x - 2)$   
 c  $(3x + 1)(x - 5)$       d  $(x - 20)(x + 20)$   
 3 a





- 4 a  $x < 3$     b  $x \geq 9$     c  $x \leq 2.5$     d  $x > -7$

**Exercise 2A**

- 1 a  $x = -1$  or  $x = -2$     b  $x = -1$  or  $x = -4$   
 c  $x = -5$  or  $x = -2$     d  $x = 3$  or  $x = -2$   
 e  $x = 3$  or  $x = 5$     f  $x = 4$  or  $x = 5$   
 g  $x = 6$  or  $x = -1$     h  $x = 6$  or  $x = -2$   
 2 a  $x = 0$  or  $x = 4$     b  $x = 0$  or  $x = 25$   
 c  $x = 0$  or  $x = 2$     d  $x = 0$  or  $x = 6$   
 e  $x = -\frac{1}{2}$  or  $x = -3$     f  $x = -\frac{1}{3}$  or  $x = \frac{3}{2}$   
 g  $x = -\frac{2}{3}$  or  $x = \frac{3}{2}$     h  $x = \frac{3}{2}$  or  $x = \frac{5}{2}$   
 3 a  $x = \frac{1}{3}$  or  $x = -2$     b  $x = 3$  or  $x = 0$   
 c  $x = 13$  or  $x = 1$     d  $x = 2$  or  $x = -2$   
 e  $x = \pm\sqrt{\frac{5}{3}}$     f  $x = 3 \pm \sqrt{13}$   
 g  $x = \frac{1 \pm \sqrt{11}}{3}$     h  $x = 1$  or  $x = -\frac{7}{6}$   
 i  $x = -\frac{1}{2}$  or  $x = \frac{7}{3}$     j  $x = 0$  or  $x = -\frac{11}{6}$   
 4  $x = 4$   
 5  $x = -1$  or  $x = -\frac{2}{25}$

**Exercise 2B**

- 1 a  $x = \frac{1}{2}(-3 \pm \sqrt{5})$     b  $x = \frac{1}{2}(3 \pm \sqrt{17})$   
 c  $x = -3 \pm \sqrt{3}$     d  $x = \frac{1}{2}(5 \pm \sqrt{33})$   
 e  $x = \frac{1}{3}(-5 \pm \sqrt{31})$     f  $x = \frac{1}{2}(1 \pm \sqrt{2})$   
 g  $x = 2$  or  $x = -\frac{1}{4}$     h  $x = \frac{1}{11}(-1 \pm \sqrt{78})$   
 2 a  $x = -0.586$  or  $x = -3.41$     b  $x = 7.87$  or  $x = 0.127$   
 c  $x = 0.765$  or  $x = -11.8$     d  $x = 8.91$  or  $x = -1.91$   
 e  $x = 0.105$  or  $x = -1.90$     f  $x = 3.84$  or  $x = -2.34$   
 g  $x = 4.77$  or  $x = 0.558$     h  $x = 4.89$  or  $x = -1.23$   
 3 a  $x = -6$  or  $x = -2$     b  $x = 1.09$  or  $x = -10.1$   
 c  $x = 9.11$  or  $x = -0.110$     d  $x = -\frac{1}{2}$  or  $x = -2$   
 e  $x = 1$  or  $x = -9$     f  $x = 1$   
 g  $x = 4.68$  or  $x = -1.18$     h  $x = 3$  or  $x = 5$   
 4 Area =  $\frac{1}{2}(2x)(x + (x+10)) = 50 \text{ m}^2$   
 So  $x^2 + 5x - 25 = 0$   
 Using the quadratic formula:  
 $x = \frac{1}{2}(-5 \pm 5\sqrt{5})$   
 Height =  $2x = 5(\sqrt{5} - 1) \text{ m}$

**Challenge**

$x = 13$

**Exercise 2C**

- 1 a  $(x + 2)^2 - 4$     b  $(x - 3)^2 - 9$   
 c  $(x - 8)^2 - 64$     d  $(x + \frac{1}{2})^2 - \frac{1}{4}$   
 e  $(x - 7)^2 - 49$   
 2 a  $2(x + 4)^2 - 32$     b  $3(x - 4)^2 - 48$   
 c  $5(x + 2)^2 - 20$     d  $2(x - \frac{5}{4})^2 - \frac{25}{8}$   
 e  $-2(x - 2)^2 + 8$   
 3 a  $2(x + 2)^2 - 7$     b  $5(x - \frac{3}{2})^2 - \frac{33}{4}$   
 c  $3(x + \frac{1}{3})^2 - \frac{4}{3}$     d  $-4(x + 2)^2 + 26$   
 e  $-8(x - \frac{1}{8})^2 + \frac{81}{8}$

- 4  $a = \frac{3}{2}, b = \frac{15}{4}$   
 5  $A = 6, B = 0.04, C = -10$

**Exercise 2D**

- 1 a  $x = -3 \pm 2\sqrt{2}$     b  $x = -6 \pm \sqrt{33}$   
 c  $x = -2 \pm \sqrt{6}$     d  $x = 5 \pm \sqrt{30}$   
 2 a  $x = \frac{1}{2}(-3 \pm \sqrt{15})$     b  $x = \frac{1}{5}(-4 \pm \sqrt{26})$   
 c  $x = \frac{1}{8}(1 \pm \sqrt{129})$     d  $x = \frac{1}{2}(-3 \pm \sqrt{39})$   
 3 a  $p = -7, q = -48$   
 b  $(x - 7)^2 = 48$   
 $x = 7 \pm \sqrt{48} = 7 \pm 4\sqrt{3}$   
 $r = 7, s = 4$   
 4  $x^2 + 2bx + c = (x + b)^2 - b^2 + c$   
 $(x + b)^2 = b^2 - c$   
 $x = -b \pm \sqrt{b^2 - c}$


**Challenge**

- a  $ax^2 + 2bx + c = 0$     b  $ax^2 + bx + c = 0$   
 $x^2 + \frac{2b}{a}x + \frac{c}{a} = 0$      $x^2 + \frac{b}{a}x + \frac{c}{a} = 0$   
 $(x + \frac{b}{a})^2 - \frac{b^2}{a^2} + \frac{c}{a} = 0$      $(x + \frac{b}{2a})^2 - \frac{b^2}{4a^2} + \frac{c}{a} = 0$   
 $(x + \frac{b}{a})^2 = \frac{b^2 - ac}{a^2}$      $(x + \frac{b}{2a})^2 = \frac{b^2 - 4ac}{4a^2}$   
 $x = -\frac{b}{a} \pm \sqrt{\frac{b^2 - ac}{a^2}}$      $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

**Exercise 2E**


- 1 a 8    b 7    c 3    d 10.5    e 0  
 f 0    g 25    h 2    i 7  
 2  $\alpha = 4$  or  $\alpha = -2$   
 3 a  $\frac{2}{3}$     b 2 and -9    c -10 and 4  
 d 12 and -12    e 0, -5 and -7    f 0, 3 and -8  
 4  $x = 3$  and  $x = 2$   
 5  $x = 0, x = 2.5$  and 6  
 6 a  $(x - 1)^2 + 1$   
 $p = -1, q = 1$   
 b Squared terms are always  $\geq 0$ , so the minimum value is  $0 + 1 = 1$   
 7 a -2 and -1    b 2, -2,  $2\sqrt{2}$  and  $-2\sqrt{2}$   
 c -1 and  $\frac{1}{3}$     d  $\frac{1}{2}$  and 1  
 e 4 and 25    f 8 and -27  
 8 a  $(3^x - 27)(3^x - 1)$     b 0 and 3

**Exercise 2F**




Turning point: (3, -1)  
 Line of symmetry:  $x = 3$







Turning point:  $(-1, -16)$   
Line of symmetry:  $x = -1$




Turning point:  $(0, 25)$   
Line of symmetry:  $x = 0$




Turning point:  $(-\frac{3}{2}, -\frac{1}{4})$   
Line of symmetry:  $x = -\frac{3}{2}$




Turning point:  $(3, 16)$   
Line of symmetry:  $x = 3$




Turning point:  $(-1, 8)$   
Line of symmetry:  $x = -1$




Turning point:  $(-\frac{7}{4}, -\frac{169}{8})$   
Line of symmetry:  $x = -\frac{7}{4}$



Turning point:  $(\frac{19}{12}, -\frac{121}{24})$   
Line of symmetry:  $x = \frac{19}{12}$



Turning point:  $(-\frac{7}{4}, \frac{81}{8})$   
Line of symmetry:  $x = -\frac{7}{4}$



Turning point:  $(-0.2, 0)$   
Line of symmetry:  $x = -0.2$

- 2 **a**  $a = 1, b = -8, c = 15$   
**b**  $a = -1, b = 3, c = 10$   
**c**  $a = 2, b = 0, c = -18$   
**d**  $a = \frac{1}{4}, b = -\frac{3}{4}, c = -1$
- 3 **a**  $a = 3, b = -30, c = 72$

**Exercise 2G**

- 1 **a** **i** 52                      **ii** -23                      **iii** 37  
          **iv** 0                        **v** -44
- b** **i**  $h(x)$                       **ii**  $f(x)$                       **iii**  $k(x)$   
          **iv**  $j(x)$                       **v**  $g(x)$
- 2  $k < 9$
- 3  $t = \frac{9}{8}$
- 4  $s = 4$
- 5  $k > \frac{4}{3}$
- 6 **a**  $p = 6$                       **b**  $x = -9$
- 7 **a**  $k^2 + 16$   
**b**  $k^2$  is always positive so  $k^2 + 16 > 0$

**Challenge**

- a** Need  $b^2 > 4ac$ . If  $a, c > 0$  or  $a, c < 0$ , choose  $b$  such that  $b > \sqrt{4ac}$ . If  $a > 0$  and  $c < 0$  (or vice versa), then  $4ac < 0$ , so  $4ac < b^2$  for all  $b$ .
- b** Not if one of  $a$  or  $c$  are negative as this would require  $b$  to be the square root of a negative number. Possible if both negative or both positive.





**Exercise 2H**





- 1 a The height of the bridge above ground level  
 b  $x = 1103$  and  $x = -1103$   
 c 2206 m
- 2 a 21.8 mph and 75.7 mph  
 b  $A = 39.77, B = 0.01, C = 48.75$   
 c 48.75 mph  
 d -11 mpg; a negative answer is impossible so this model is not valid for very high speeds
- 3 a 6 tonnes  
 b 39.6 kilograms per hectare.
- 4 a  $M = 40\,000$   
 b  $r = 400\,000 - 1000(p - 20)^2$   
 $A = 400\,000, B = 1000, C = 20$   
 c £20

**Challenge**

- a  $a = 0.01, b = 0.3, c = -4$   
 b 36.2 mph

**Mixed exercise**

- 1 a  $y = -1$  or  $-2$                       b  $x = \frac{2}{3}$  or  $x = -5$   
 c  $x = -\frac{1}{5}$  or 3                              d  $\frac{5 \pm \sqrt{7}}{2}$



- 3 a  $k = 1$                       b  $x = 3$  and  $x = -2$
- 4 a  $k = 0.0902$  or  $k = -11.1$   
 b  $t = 2.28$  or  $t = 0.219$   
 c  $x = -2.30$  or  $x = 1.30$   
 d  $x = 0.839$  or  $x = -0.239$
- 5 a  $(x + 6)^2 - 45; p = 1, q = 6, r = -45$   
 b  $5(x - 4)^2 - 67; p = 5, q = -4, r = -67$   
 c  $-2(x - 2)^2 + 8; p = -2, q = -2, r = 8$   
 d  $2(x - \frac{1}{2})^2 - \frac{3}{2}; p = 2, q = -\frac{1}{2}, r = -\frac{3}{2}$

- 6  $k = \frac{1}{5}$
- 7 a  $p = 3, q = 2, r = -7$                       b  $-2 \pm \sqrt{\frac{7}{3}}$
- 8 a  $f(x) = (2x - 16)(2x - 4)$                       b 4 and 2
- 9  $1 \pm \sqrt{13}$
- 10  $x = -5$  or  $x = 4$
- 11 a 10 m                      b 1.28 s  
 c  $h(t) = 10.625 - 10(t - 0.25)^2$   
 $A = 10.625, B = 10, C = 0.25$   
 d 10.625 m at 0.25 s
- 12 a  $16k^2 + 4$   
 b  $k^2 \geq 0$  for all  $k$ , so  $16k^2 + 4 > 0$   
 c When  $k = 0, f(x) = 2x + 1$ ; this is a linear function with only one root
- 13 1, -1, 2 and -2
- 14 a  $H = 10$   
 b  $r = 1322.5 - 10(p - 11.5)^2$   
 $A = 1322.5, B = 10, C = 11.5$   
 c Old revenue is  $80 \times £15 = £1200$ ; new revenue is £1322.50; difference is £122.50. The best selling price of a cushion is £11.50.

**Challenge**

- a  $\frac{a+b}{a} = \frac{a}{b}$   
 $a^2 - ba - b^2 = 0$   
 Using quadratic formula:  $a = \frac{b + \sqrt{5b^2}}{2}$   
 So  $a : b$  is  $\frac{b + \sqrt{5b^2}}{2} : b$   
 Dividing by  $b: \frac{1 + \sqrt{5}}{2} : 1$
- b Let  $x = \sqrt{1 + \sqrt{1 + \sqrt{1 + \sqrt{1 + \dots}}}}$   
 So  $x = \sqrt{1 + x} \Rightarrow x^2 - x - 1 = 0$   
 Using quadratic formula:  $x = \frac{1 + \sqrt{5}}{2}$

**CHAPTER 3**

**Prior knowledge check**

- 1 a  $A \cap B = \{1, 2, 4\}$                       b  $(A \cup B)' = \{7, 9, 11, 13\}$   
 2 a  $5\sqrt{3}$                                       c  $\sqrt{5} + 2\sqrt{2}$   
 3 a graph ii                      b graph iii                      c graph i

**Exercise 3A**

- 1 a  $x = 4, y = 2$                       b  $x = 1, y = 3$   
 c  $x = 2, y = -2$                       d  $x = 4\frac{1}{2}, y = -3$   
 e  $x = -\frac{2}{3}, y = 2$                       f  $x = 3, y = 3$
- 2 a  $x = 5, y = 2$                       b  $x = 5\frac{1}{2}, y = -6$   
 c  $x = 1, y = -4$                       d  $x = 1\frac{3}{4}, y = \frac{1}{4}$
- 3 a  $x = -1, y = 1$                       b  $x = 4, y = -4$   
 c  $x = 0.5, y = -2.5$
- 4 a  $3x + ky = 8$  (1);  $x - 2ky = 5$  (2)  
 $(1) \times 2: 6x + 2ky = 16$  (3)  
 $(2) + (3) 7x = 21$  so  $x = 3$   
 b -2
- 5  $p = 3, q = 1$

**Exercise 3B**


- 1 a  $x = 5, y = 6$  or  $x = 6, y = 5$   
 b  $x = 0, y = 1$  or  $x = \frac{4}{5}, y = -\frac{3}{5}$   
 c  $x = -1, y = -3$  or  $x = 1, y = 3$

- d  $a = 1, b = 5$  or  $a = 3, b = -1$   
 e  $u = 1\frac{1}{2}, v = 4$  or  $u = 2, v = 3$   
 f  $x = -1\frac{1}{2}, y = 5\frac{3}{4}$  or  $x = 3, y = -1$   
 2 a  $x = 3, y = \frac{1}{2}$  or  $x = 6\frac{1}{3}, y = -2\frac{5}{6}$   
 b  $x = 4\frac{1}{2}, y = 4\frac{1}{2}$  or  $x = 6, y = 3$   
 c  $x = -19, y = -15$  or  $x = 6, y = 5$   
 3 a  $x = 3 + \sqrt{13}, y = -3 + \sqrt{13}$  or  $x = 3 - \sqrt{13}, y = -3 - \sqrt{13}$   
 b  $x = 2 - 3\sqrt{5}, y = 3 + 2\sqrt{5}$  or  $x = 2 + 3\sqrt{5}, y = 3 - 2\sqrt{5}$   
 4  $x = -5, y = 8$  or  $x = 2, y = 1$   
 5 a  $3x^2 + x(2 - 4x) + 11 = 0$   
 $3x^2 + 2x - 4x^2 + 11 = 0$   
 $x^2 - 2x - 11 = 0$   
 b  $x = 1 + 2\sqrt{3}, y = -2 - 8\sqrt{3}$   
 $x = 1 - 2\sqrt{3}, y = -2 + 8\sqrt{3}$   
 6 a  $k = 3, p = -2$   
 b  $x = -6, y = -23$  or  $x = 1, y = -2$


**Challenge**

$y = x + k$   
 $x^2 + (x + k)^2 = 4$   
 $x^2 + x^2 + 2kx + k^2 - 4 = 0$   
 $2x^2 + 2kx + k^2 - 4 = 0$  for one solution  $b^2 - 4ac = 0$   
 $4k^2 - 4 \times 2(k^2 - 4) = 0$   
 $4k^2 - 8k^2 + 32 = 0$   $4k^2 = 32$   $k^2 = 8$   $k = \pm 2\sqrt{2}$


**Exercise 3C**




ii (2, 1)




ii (3, -1)




ii (-0.5, 0.5)



b (3.5, 9) and (-1.5, 4)



b (-1, 8) and (3, 0)



b (6, 16) and (1, 1)

5 (-11, -15) and (3, -1)

6  $(-1\frac{1}{6}, -4\frac{1}{2})$  and (2, 5)

7 a 2 points      b 1 point      c 0 points

8 a  $y = 2x - 1$   
 $x^2 + 4k(2x - 1) + 5k = 0$   
 $x^2 + 8kx - 4k + 5k = 0$        $x^2 + 8kx + k = 0$

b  $k = \frac{1}{16}$       c  $x = -\frac{1}{4}, y = -\frac{3}{2}$

9 If swimmer reaches the bottom of the pool

$0.5x^2 - 3x = 0.3x - 6$

$0.5x^2 - 3.3x + 6 = 0$

$b^2 - 4ac = (-3.3)^2 - 4 \times 0.5 \times 6 = -1.11$

negative so no points of intersection and diver does not reach the bottom of the pool

**Exercise 3D**

1 a  $x < 4$       b  $x \geq 7$       c  $x > 2\frac{1}{2}$       d  $x \leq -3$

e  $x < 11$       f  $x < 2\frac{3}{5}$       g  $x > -12$       h  $x < 1$

i  $x \leq 8$       j  $x > 1\frac{1}{7}$

2 a  $x \geq 3$       b  $x < 1$       c  $x \leq -3\frac{1}{4}$       d  $x < 18$

e  $x > 3$       f  $x \geq 4\frac{2}{5}$       g  $x < 4$       h  $x > -7$

i  $x \leq -\frac{1}{2}$       j  $x \geq \frac{3}{4}$       k  $x \geq -\frac{10}{3}$       l  $x \geq \frac{9}{11}$

3 a  $\{x: x > 2\frac{1}{2}\}$       b  $\{x: 2 < x < 4\}$

c  $\{x: 2\frac{1}{2} < x < 3\}$       d No values

e  $x = 4$

f  $\{x: x < 1.2\} \cup \{x: x > 2.2\}$

g  $\{x: x \leq -\frac{2}{3}\} \cup \{x: x \geq \frac{3}{2}\}$

**Challenge**

$p = -1, q = 4, r = 6$

**Exercise 3E**

1 a  $3 < x < 8$       b  $-4 < x < 3$

c  $x < -2, x > 5$       d  $x \leq -4, x \geq -3$

e  $-\frac{1}{2} < x < 7$       f  $x < -2, x > 2\frac{1}{2}$

g  $\frac{1}{2} \leq x \leq 1\frac{1}{2}$       h  $x < \frac{1}{3}, x > 2$

i  $-3 < x < 3$       j  $x < -2\frac{1}{2}, x > \frac{2}{3}$

k  $x < 0, x > 5$       l  $-1\frac{1}{2} \leq x \leq 0$

2 a  $-5 < x < 2$       b  $x < -1, x > 1$

c  $\frac{1}{2} < x < 1$       d  $-3 < x < \frac{1}{4}$

3 a  $\{x: 2 < x < 4\}$       b  $\{x: x > 3\}$


c  $\{x: -\frac{1}{4} < x < 0\}$       d No values




- e  $\{x: -5 < x < -3\} \cup \{x: x > 4\}$   
 f  $\{x: -1 < x < 1\} \cup \{x: 2 < x < 3\}$
- 4 a  $x < 0$  or  $x > 2$       b  $x < 0$  or  $x > 0.8$   
 c  $x < -1$  or  $x > 0$       d  $x < 0$  or  $x > 0.5$   
 e  $x < -\frac{1}{5}$  or  $x > \frac{1}{5}$       f  $x \leq -\frac{2}{3}$  or  $x \geq 3$
- 5 a  $-2 < k < 6$       b  $p \leq -8$  or  $p \geq 0$
- 6  $\{x: x < -2\} \cup \{x: x > 7\}$
- 7 a  $\{x: x < \frac{2}{3}\}$       b  $\{x: -\frac{1}{2} < x < 3\}$   
 c  $\{x: -\frac{1}{2} < x < \frac{2}{3}\}$
- 8  $x < 3$  or  $x > 5.5$
- 9 No real roots  $b^2 - 4ac < 0$        $(-2k)^2 - 4 \times k \times 3 < 0$   
 $4k^2 - 12k = 0$  when  $k = 0$  and  $k = 3$   
 solution  $0 \leq k < 3$   
 note when  $k = 0$  equation gives  $3 = 0$

**Exercise 3F**


- 1 a  $P(3.2, -1.8)$       b  $x < 3.2$




- ii  $(4, 5)$       iii  $x \leq 4$




- ii  $(-3, 23)$       iii  $x \geq -3$




- ii  $(-2, 9), (0, 5)$       iii  $-2 \leq x \leq 0$



- ii  $(-5, -22), (3, -6)$       iii  $x \leq -5$  or  $x \geq 3$



- ii  $(-2, -1), (9, 76)$       iii  $-2 \leq x \leq 9$








- ii  $(-5, -18), (3, -2)$       iii  $x \leq -5$  or  $x \geq 3$
- 3 a  $-1 < x < 2$       b  $0.5 < x < 3$   
 c  $x < 0.5$  or  $x > 3$       d  $x < 0$  or  $x > 2$   
 e  $1 < x < 3$       f  $x < -1$  or  $x > -0.75$


**Challenge**

- a  $(-1.5, -3.75), (6, 0)$   
 b  $\{x: -1.5 < x < 6\}$

**Exercise 3G**



- 6 a  $(1, 6), (3, 4), (1, 2)$   
 b  $x \geq 1, y \leq 7 - x, y \geq x + 1$   
 7  $y < 2 - 5x - x^2, 2x + y \geq 0, x + y \leq 4$




- b  $(-\frac{7}{6}, \frac{17}{6}), (2, 6), (2, -1), (-0.4, 1)$   
 c  $(-0.4, 1)$       d  $\frac{941}{60}$





**Mixed exercise**

- 1 a  $4kx - 2y = 8$   
 $4kx + 3y = -2$   
 $-5y = 10$   
 $y = -2$
- b  $x = \frac{1}{k}$
- 2  $x = -4, y = 3\frac{1}{2}$
- 3 a Substitute  $x = 1 + 2y$  into  $3xy - y^2 = 8$   
b  $(3, 1)$  and  $(-\frac{11}{5}, -\frac{8}{5})$
- 4 a Substitute  $y = 2 - x$  into  $x^2 + xy - y^2 = 0$   
b  $x = 3 \pm \sqrt{6}, y = -1 \pm \sqrt{6}$
- 5 a  $3^x = (3^2)^{y-1} = 3^{2y-2} \Rightarrow x = 2y - 2$   
b  $x = 4, y = 3$  and  $x = -\frac{2}{3}, y = -\frac{1}{3}$
- 6  $x = -1\frac{1}{2}, y = 2\frac{1}{4}$  and  $x = 4, y = -\frac{1}{2}$
- 7 a  $k = -2$                                 b  $(-1, 2)$
- 8 Yes, the ball will hit the ceiling
- 9 a  $\{x: x > 10\frac{1}{2}\}$                                 b  $\{x: x < -2\} \cup \{x: x > 7\}$
- 10  $3 < x < 4$
- 11 a  $x = -5, x = 4$                                 b  $\{x: x < -5\} \cup \{x: x > 4\}$
- 12 a  $x < 2\frac{1}{2}$                                         b  $\frac{1}{2} < x < 5$   
c  $0 < x < 4$                                         d  $\frac{1}{2} < x < 2\frac{1}{2}$
- 13  $1 \leq x \leq 8$
- 14  $k \leq 3\frac{1}{5}$
- 15  $b^2 < 4ac$  so  $16k^2 < -40k$   
 $8k(2k + 5) < 0$  so  $-\frac{5}{2} < k < 0$



- b  $(-7, 20), (3, 0)$                                 c  $x < -7, x > 3$

17  $\frac{1}{4}(-1 - \sqrt{185}) < x < \frac{1}{4}(-1 + \sqrt{185})$



b  $\frac{9}{2}$

**Challenge**

- 1  $0 < x < 1.6$                                 2  $-2 < k < 7$

**CHAPTER 4**


**Prior knowledge check**

- 1 a  $(x + 5)(x + 1)$                                 b  $(x - 3)(x - 1)$
- 2 a                                b

3 a

$x$	-2	-1.5	-1	-0.5	0
$y$	-12	-6.875	-4	-2.625	-2

$x$	0.5	1	1.5	2
$y$	-1.375	0	2.875	8













- 4 a  $x = 2, y = 4$                                 b  $x = 2, y = 5$

**Exercise 4A**













- 1 a                                b
- c                                d
- e                                f
- g                                h









**i**  $y = x(x - 3)(x + 3)$

**j**  $y = x^2(x - 9)$





**3 a**  $y = x(x + 2)(x - 1)$









**b**  $y = x(x + 4)(x + 1)$



**c**  $y = x(x + 1)^2$






**d**  $y = x(x + 1)(3 - x)$







- 5 a  $b = 4, c = 1, d = -6$   
 6  $a = \frac{1}{3}, b = -\frac{4}{3}, c = \frac{1}{3}, d = 2$   
 7 a  $x(x^2 - 12x + 32)$






- b  $(0, -6)$   
 b  $x(x - 8)(x - 4)$



**Exercise 4B**



- 3 a  $(0, 12)$   
 b  $b = -2, c = -7, d = 8, e = 12$



**Challenge**


$a = \frac{1}{3}, b = -\frac{4}{3}, c = -\frac{2}{3}, d = 4, e = 3$






**Exercise 4C**


1 a




b




c




d




e




2 a



b




c



**Exercise 4D**


1 a i



ii 3

iii  $x^2 = x(x^2 - 1)$


b i



ii 1

iii  $x(x + 2) = -\frac{3}{x}$


c i



ii 3

iii  $x^2 = (x + 1)(x - 1)^2$


d i



ii 2

iii  $x^2(1 - x) = \frac{2}{x}$


e i



ii 1


iii  $x(x - 4) = \frac{1}{x}$

f i




ii 3


iii  $x(x - 4) = -\frac{1}{x}$




**ii** 1      **iii**  $x(x-4) = (x-2)^3$




**ii** 2      **iii**  $-x^3 = -\frac{2}{x}$




**ii** 2      **iii**  $-x^3 = x^2$




**ii** 3      **iii**  $-x^3 = -x(x+2)$




**ii** 2      **iii**  $x(x-1)(x+2)^2 = 4$




**ii** 1      **iii**  $x^3 = x^2(x+1)^2$




**b** Only 2 intersections



**b** Only 1 intersection




**b** Graphs do not intersect




**b** 2; the graphs cross in two places so there are two solutions.







- b 3  
c Expand brackets and rearrange.  
d  $(-2, 1)$ ,  $(-1, 4)$ ,  $(\frac{2}{3}, 9)$




- b  $(0, 0)$ ;  $(-2, -12)$ ;  $(5, 30)$




- b  $(0, 2)$ ;  $(-3, -40)$ ;  $(5, 72)$



- b  $(0, -8)$ ;  $(1, -9)$ ;  $(-4, -24)$








- b Graphs do not intersect.  
c  $a < -\frac{7}{16}$






- b 2

Exercise 4E












7 a (6, -1) b (4, 2)


8  $y = \frac{1}{x-4}$




c  $f(x+2) = (x+1)(x+4)$ ; (0, 4)  
 $f(x)+2 = (x+1)(x+2)+2$ ; (0, 0)



c  $f(x+1) = -x(x+1)^2$ ; (0, 0)



b -1 or 1




b -2, -3 or -5

**Challenge**


1 (3, 2)
















2 a (-7, -12) b  $f(x-2)+1$





c  $f(x+2) = (x+2)(x-2)$ ; (2, 0); (-2, 0)  
 $f(x)+4 = (x-2)^2$ ; (2, 0)

**Exercise 4F**








7 a (1, -3)      b (2, -12)

8 (-4, 8)











b 2 and 3

**Challenge**


1 (2, -2)

2  $\frac{1}{4}f(\frac{1}{2}x)$


**Exercise 4G**




2 a  $y = 4, x = 1, (0, 2)$




b  $y = 2, x = 0, (-1, 0)$




c  $y = 4, x = 1, (0, 0)$



d  $y = 0, x = 1, (0, -2)$




e  $y = 2, x = \frac{1}{2}, (0, 0)$







f  $y = 2, x = 2, (0, 0)$




g  $y = 1, x = 1, (0, 0)$




h  $y = -2, x = 1, (0, 0)$




3 a  $A(-2, -6), B(0, 0), C(2, -3), D(6, 0)$




b  $A(-4, 0), B(-2, 6), C(0, 3), D(4, 6)$




c  $A(-2, -6), B(-1, 0), C(0, -3), D(2, 0)$




d  $A(-8, -6), B(-6, 0), C(-4, -3), D(0, 0)$




e  $A(-4, -3), B(-2, 3), C(0, 0), D(4, 3)$




f  $A(-4, -18), B(-2, 0), C(0, -9), D(4, 0)$




g  $A(-4, -2), B(-2, 0), C(0, -1), D(4, 0)$




h  $A(-16, -6), B(-8, 0), C(0, -3), D(16, 0)$




i  $A(-4, 6), B(-2, 0), C(0, 3), D(4, 0)$




j  $A(4, -6), B(2, 0), C(0, -3), D(-4, 0)$




4 a i  $x = -2, y = 0, (0, 2)$




ii  $x = -1, y = 0, (0, 1)$




iii  $x = 0, y = 0$




iv  $x = -2, y = -1, (0, 0)$



v  $x = 2, y = 0, (0, 1)$



vi  $x = -2, y = 0, (0, -1)$




b  $f(x) = \frac{2}{x+1}$


5 a  $\frac{1}{2}$

b i (6, 1) ii (2, 3) iii (2, -3.5)


6 a  $A(-1, -2) B(0, 0) C(1, 0) D(2, -2)$




b  $A(-1, 0) B(0, 4) C(1, 4) D(2, 0)$




c  $A(-1, 3) B(0, 5) C(1, 5) D(2, 3)$



d  $A(-1, 0) B(0, \frac{2}{3}) C(1, \frac{2}{3}) D(2, 0)$




e  $A(-1, 0.5) B(0, 1.5) C(1, 1.5) D(2, 0.5)$




**Mixed exercise**

1 a  $y = x^2(x - 2)$




b  $x = 0, -1, 2$ ; points  $(0, 0), (2, 0), (-1, -3)$




b  $A(-3, -2), B(2, 3)$

c  $y = x^2 + 2x - 5$


3 a  $A(\frac{3}{2}, 4)$






b  $A(3, 2)$




c  $A(3, 2)$   
 $B(0, -2)$   
 $y = 0$  is asymptote





d  $A(0, 4)$





4 a  $x = -1$  at A,  $x = 3$  at B



6 a  $(0, 2)$



7 a  $y = x^2 - 4x + 3$



8 a  $(0, 2)$       b  $-2$       c  $-1, 1, 2$


9 a i  $(\frac{4}{3}, 3)$       ii  $(4, 6)$       iii  $(9, 3)$

iv  $(4, -3)$       v  $(4, -\frac{1}{2})$

b  $f(2x), f(x + 2)$


c i  $f(x - 4) + 3$       ii  $2f(\frac{1}{2}x)$

10 a




b 1; only one intersection of the two curves

11 a  $x(x - 3)^2$




c -4 and -7


12 a  $y = x(x - 2)^2$




b  $y = x(x - 2)^2 + x$



13 a Asymptotes at  $x = 0$  and  $y = -2$



b  $(\frac{1}{2}, 0)$



d Asymptotes at  $y = 0$  and  $x = -3$ ; intersection at  $(0, \frac{1}{3})$


Challenge

$(6 - c, -4 - d)$


Review exercise 1

- 1 a 2
- 2 a 625
- 3 a  $4\sqrt{5}$
- 4 a 13
- 5 a  $1 + 2\sqrt{k}$
- 6 a  $25x^{-4}$
- 7  $8 + 8\sqrt{2}$
- 8  $1 - 2\sqrt{2}$
- 9 a  $(x - 8)(x - 2)$
- 10 a  $a = -4, b = -45$
- 11 4.19 (3 s.f.)
- 12 a The height of the athlete's shoulder is 1.7 m
- 13 a  $(x - 3)^2 + 9$
- 14 a  $k = 2$


- b  $\frac{1}{4}$
- b  $\frac{4}{3}x^{\frac{2}{3}}$
- b  $21 - 8\sqrt{5}$
- b  $8 - 2\sqrt{3}$
- b  $1 + 6\sqrt{k}$
- b  $x^2$
- b  $y = 1, y = \frac{1}{3}$
- b  $x = 4 \pm 3\sqrt{5}$



- 15 a  $x^3(x^3 - 8)(x^3 + 1)$
- 16 a  $a = 5, b = 11$
- b discriminant  $< 0$  so no real roots
- c  $k = 25$



- 17 a  $a = 1, b = 2$
- b



- c discriminant = -8
- d  $-2\sqrt{3}k < 2\sqrt{3}$
- 18 a  $x^2 + 4x - 8 = 0$
- b  $x = -2 \pm 2\sqrt{3}, y = -6 \pm 2\sqrt{3}$
- 19 a  $x > \frac{1}{4}$
- b  $x < \frac{1}{2}$  or  $x > 3$
- c  $\frac{1}{4} < x < \frac{1}{2}$  or  $x > 3$
- 20  $-2(x + 1) = x^2 - 5x + 2$
- $x^2 - 3x + 4 = 0$

The discriminant of this is  $-7 < 0$ , so no real solutions.





21 a  $x = \frac{7}{2}, y = -2, x = -3, y = 11$


b  $x < -3$  or  $x > 3\frac{1}{2}$

22 a Different real roots, discriminant  $> 0$   
so  $k^2 - 4k - 12 > 0$




b  $k < -2$  or  $k > 6$

23  $-7 < x < 2$


24




25 a  $x(x-2)(x+2)$




(2, 0) (4, 0) and (3, 2)




(1, 0) (2, 0) and  $(1\frac{1}{2}, -2)$




(0, 0) and (3, 0)



(1, 0) (4, 0) and (0, 6)




(2, 0) (8, 0) and (0, 3)




Asymptotes:  $y = 3$  and  $x = 0$

b  $(-\frac{1}{3}, 0)$


29 a 0.438, 1, 4, 4.56



30 a (6, 8)    b (9, -8)    c (6, -4)



b 1



b  $-\frac{1}{2}, \frac{1}{2}$

**Challenge**

1 a  $x = 1, x = 9$     b 0, 2

2  $\sqrt{2}$  cm,  $3\sqrt{2}$  cm

3  $3x^3 + x^2 - x = 2x(x-1)(x+1)$

$3x^3 + x^2 - x = 2x^3 - 2x$

$x^3 + x^2 + x = 0$

$x(x^2 + x + 1) = 0$

The discriminant of the bracket is  $-3 < 0$  so this contributes no real solutions.

The only solution is when  $x = 0$  at (0, 0).

4 -3, 3

**CHAPTER 5**

**Prior knowledge check**

- 1 a  $(-2, -1)$     b  $(\frac{9}{19}, \frac{26}{19})$     c  $(7, 3)$   
 2 a  $4\sqrt{5}$     b  $10\sqrt{2}$     c  $5\sqrt{5}$   
 3 a  $y = 5 - 2x$     b  $y = \frac{2}{5}x - \frac{9}{5}$     c  $y = \frac{3}{12}x + \frac{12}{7}$

**Exercise 5A**

- 1 a  $\frac{1}{2}$     b  $\frac{1}{6}$     c  $-\frac{3}{5}$     d 2  
 e -1    f  $\frac{1}{2}$     g  $\frac{1}{2}$     h 8  
 i  $\frac{2}{3}$     j -4    k  $-\frac{1}{3}$     l  $-\frac{1}{2}$   
 m 1    n  $\frac{q^2 - p^2}{q - p} = q + p$   
 2 7  
 3 12  
 4  $4\frac{1}{3}$   
 5  $2\frac{1}{4}$   
 6  $\frac{1}{4}$   
 7 26  
 8 -5  
 9 Gradient of  $AB =$  gradient of  $BC = 0.5$ ; point  $B$  is common  
 10 Gradient of  $AB =$  gradient of  $BC = -0.5$ ; point  $B$  is common

**Exercise 5B**

- 1 a -2    b -1    c 3    d  $\frac{1}{3}$   
 e  $-\frac{2}{3}$     f  $\frac{5}{4}$     g  $\frac{1}{2}$     h 2  
 i  $\frac{1}{2}$     j  $\frac{1}{2}$     k -2    l  $-\frac{3}{2}$   
 2 a 4    b -5    c  $-\frac{2}{3}$     d 0  
 e  $\frac{7}{5}$     f 2    g 2    h -2  
 i 9    j -3    k  $\frac{3}{2}$     l  $-\frac{1}{2}$   
 3 a  $4x - y + 3 = 0$     b  $3x - y - 2 = 0$   
 c  $6x + y - 7 = 0$     d  $4x - 5y - 30 = 0$   
 e  $5x - 3y + 6 = 0$     f  $7x - 3y = 0$   
 g  $14x - 7y - 4 = 0$     h  $27x + 9y - 2 = 0$   
 i  $18x + 3y + 2 = 0$     j  $2x + 6y - 3 = 0$   
 k  $4x - 6y + 5 = 0$     l  $6x - 10y + 5 = 0$   
 4  $(3, 0)$   
 5  $(0, 0)$   
 6  $(0, 5), (-4, 0)$   
 7 a  $\frac{1}{3}$     b  $x - 3y + 15 = 0$   
 8 a  $-\frac{2}{5}$     b  $2x + 5y - 10 = 0$   
 9  $ax + by + c = 0$   
 $by = -ax - c$   
 $y = (-\frac{a}{b})x - (\frac{c}{b})$   
 10  $a = 6, c = 10$   
 11  $P(3, 0)$   
 12 a -16    b -27

**Challenge**

Gradient =  $-\frac{a}{b}$ ;  $y$ -intercept =  $a$ . So  $y = -\frac{a}{b}x + a$

Rearrange to give  $ax + by - ab = 0$

**Exercise 5C**

- 1 a  $y = 2x + 1$     b  $y = 3x + 7$     c  $y = -x - 3$   
 d  $y = -4x - 11$     e  $y = \frac{1}{2}x + 12$     f  $y = -\frac{2}{3}x - 5$   
 g  $y = 2x$     h  $y = -\frac{1}{2}x + 2b$

- 2 a  $y = 4x - 4$     b  $y = x + 2$     c  $y = 2x + 4$   
 d  $y = 4x - 23$     e  $y = x - 4$     f  $y = \frac{1}{2}x + 1$   
 g  $y = -4x - 9$     h  $y = -8x - 33$     i  $y = \frac{6}{5}x$   
 j  $y = \frac{2}{7}x + \frac{5}{14}$   
 3  $5x + y - 37 = 0$   
 4  $y = x + 2, y = -\frac{1}{6}x - \frac{1}{3}, y = -6x + 23$   
 5  $a = 3, c = -27$   
 6  $a = -4, b = 8$

**Challenge**

- a  $m = \frac{(y_2 - y_1)}{(x_2 - x_1)}$   
 b  $y - y_1 = \frac{(y_2 - y_1)}{(x_2 - x_1)}(x - x_1)$   
 $\frac{(y - y_1)}{(y_2 - y_1)} = \frac{(x - x_1)}{(x_2 - x_1)}$   
 c  $y = \frac{3}{7}x + \frac{52}{7}$

**Exercise 5D**

- 1  $y = 3x - 6$     2  $y = 2x + 8$   
 3  $2x - 3y + 24 = 0$     4  $-\frac{1}{5}$   
 5  $(-3, 0)$     6  $(0, 1)$   
 7  $(0, 3\frac{1}{2})$     8  $y = \frac{2}{5}x + 3$   
 9  $2x + 3y - 12 = 0$     10  $\frac{8}{5}$   
 11  $y = \frac{4}{3}x - 4$     12  $6x + 15y - 10 = 0$   
 13  $y = -\frac{4}{5}x + 4$     14  $x - y + 5 = 0$   
 15  $y = -\frac{3}{8}x + \frac{1}{2}$     16  $y = 4x + 13$

**Exercise 5E**

- 1 a Parallel    b Not parallel    c Not parallel  
 2  $r: y = \frac{4}{5}x + 3.2, s: y = \frac{4}{5}x - 7$   
 Gradients equal therefore lines are parallel.  
 3 Gradient of  $AB = \frac{3}{5}$ , gradient of  $BC = -\frac{7}{2}$ , gradient of  $CD = \frac{3}{5}$ , gradient of  $AD = \frac{10}{3}$ . The quadrilateral has a pair of parallel sides, so it is a trapezium.  
 4  $y = 5x + 3$   
 5  $2x + 5y + 20 = 0$   
 6  $y = -\frac{1}{2}x + 7$   
 7  $y = \frac{2}{3}x$   
 8  $4x - y + 15 = 0$

**Exercise 5F**

- 1 a Perpendicular    b Parallel  
 c Neither    d Perpendicular  
 e Perpendicular    f Parallel  
 g Parallel    h Perpendicular  
 i Perpendicular    j Parallel  
 k Neither    l Perpendicular  
 2  $y = -\frac{1}{6}x + 1$   
 3  $y = \frac{8}{3}x - 8$   
 4  $y = -\frac{1}{3}x$   
 5  $y = -\frac{1}{3}x + \frac{13}{3}$   
 6  $y = -\frac{3}{2}x + \frac{17}{2}$   
 7  $3x + 2y - 5 = 0$   
 8  $7x - 4y + 2 = 0$




- 9  $l: y = -\frac{1}{3}x - 1, n: y = 3x + 5$ . Gradients are negative reciprocals, therefore lines perpendicular.
- 10  $AB: y = -\frac{1}{2}x + 4\frac{1}{2}, CD: y = -\frac{1}{2}x - \frac{1}{2}, AD: y = 2x + 7, BC: y = 2x - 13$ . Two pairs of parallel sides and lines with gradients 2 and  $-\frac{1}{2}$  are perpendicular, so  $ABCD$  is a rectangle.
- 11 a  $A(\frac{7}{5}, 0)$       b  $55x - 25y - 77 = 0$
- 12  $-\frac{9}{4}$

**Exercise 5G**


- 1 a 10      b 13      c 5      d  $\sqrt{5}$   
 e  $\sqrt{106}$       f  $\sqrt{113}$
- 2 Distance between  $A$  and  $B = \sqrt{50}$  and distance between  $B$  and  $C = \sqrt{50}$  so the lines are congruent.
- 3 Distance between  $P$  and  $Q = \sqrt{74}$  and distance between  $Q$  and  $R = \sqrt{73}$  so the lines are not congruent.
- 4  $x = -8$  or  $x = 6$
- 5  $y = -2$  or  $y = 16$
- 6 a Both lines have gradient 2.  
 b  $y = -\frac{1}{2}x + \frac{23}{2}$  or  $x + 2y - 23 = 0$   
 c  $(\frac{29}{5}, \frac{43}{5})$   
 d  $\frac{7\sqrt{5}}{5}$
- 7  $P(-\frac{3}{5}, \frac{29}{5})$  or  $P(3, -5)$
- 8 a  $AB = \sqrt{178}, BC = 3$  and  $AC = \sqrt{205}$ . All sides are different lengths, therefore the triangle is a scalene triangle.  
 b  $\frac{39}{2}$  or 19.5
- 9 a  $A(2, 11)$   
 b  $B(\frac{41}{4}, 0)$   
 c  $\frac{451}{8}$
- 10 a  $(\frac{5}{2}, 0)$       b  $(-5, 0)$   
 c  $(-10, -10)$       d  $\frac{75}{2}$
- 11 a  $y = \frac{1}{2}x - \frac{9}{2}$       b  $y = -2x + 8$   
 c  $T(0, 8)$       d  $RS = 2\sqrt{5}$  and  $TR = 5\sqrt{5}$   
 e 25
- 12 a  $x + 4y - 52 = 0$       b  $A(0, 13)$   
 c  $B(4, 12)$       d 26

**Exercise 5H**


- 1 a i  $k = 50$       ii  $d = 50t$   
 b i  $k = 0.3$  or  $\text{£}0.30$       ii  $C = 0.3t$   
 c i  $k = \frac{3}{5}$       ii  $p = \frac{3}{5}t$
- 2 a not linear




b linear



c not linear




3 a



- b The data forms a straight line, so a linear model is appropriate.
- c  $E = 0.12h + 45$
- d  $a = \text{£}0.12 = \text{cost of 1 kilowatt hour of electricity, } b = \text{£}45 = \text{fixed electricity costs (per month or per quarter)}$
- e  $\text{£}52.80$

4 a




- b The data does not follow a straight line. There is a definite curve to the points on the graph.

5 a

- a  $C = 350d + 5000$
- b  $a = 350 = \text{daily fee charged by the website designer. } b = 5000 = \text{initial cost charged by the website designer.}$
- c 24 days




- 6 a  $F = 1.8C + 32$  or  $F = \frac{9}{5}C + 32$   
 b  $a = 1.8$  = increase in Fahrenheit temperature when the Celsius temperature increases by  $1^\circ\text{C}$ .  
 $b = 32$  temperature in Fahrenheit when temperature in Celsius is  $0^\circ$ .  
 c  $38.5^\circ\text{C}$   
 d  $-40^\circ\text{C}$   
 7 a  $n = 750t + 17\,500$   
 b The increase in the number of homes receiving the internet will be the same each year.  
 8 a All the points lie close to the straight line shown.  
 b  $h = 4f + 69$   
 c 175 cm



b  $Q = 24, P = 17$

**Mixed exercise**

- 1 a  $y = -\frac{5}{12}x + \frac{11}{6}$                       b -22  
 2 a  $\frac{2k-2}{8-k} = \frac{1}{3}$  therefore  $7k = 14, k = 2$   
 b  $y = \frac{1}{3}x + \frac{1}{3}$   
 3 a  $L_1 = y = \frac{1}{7}x + \frac{12}{7}, L_2 = y = -x + 12$   
 b (9, 3)  
 4 a  $y = \frac{3}{2}x - \frac{3}{2}$                               b (3, 3)  
 5  $11x - 10y + 19 = 0$   
 6 a  $y = -\frac{1}{2}x + 3$                               b  $y = \frac{1}{4}x + \frac{9}{4}$   
 7 Gradient =  $\frac{3 + 4\sqrt{3} - 3\sqrt{3}}{2 + \sqrt{3} - 1} = \frac{3 + \sqrt{3}}{1 + \sqrt{3}} = \sqrt{3}$   
 $y = \sqrt{3}x + c$  and  $A(1, 3\sqrt{3})$ , so  $c = 2\sqrt{3}$   
 Equation of line is  $y = \sqrt{3}x + 2\sqrt{3}$   
 When  $y = 0, x = -2$ , so the line meets the  $x$ -axis at  $(-2, 0)$   
 8 a  $y = -3x + 14$                               b (0, 14)  
 9 a  $y = -\frac{1}{2}x + 4$                               b Students own work.  
 c (1, 1). Note: equation of line  $n: y = -\frac{1}{2}x + \frac{3}{2}$   
 10 20  
 11 a  $2x + y = 20$                               b  $y = \frac{1}{3}x + \frac{4}{3}$   
 12 a  $\frac{1}{2}$     b 6    c  $2x + y - 16 = 0$   
 d 10  
 13 a  $7x + 5y - 18 = 0$                               b  $\frac{162}{35}$



b  $(\frac{4}{3}, -\frac{1}{3})$                               c  $12x - 3y - 17 = 0$

- 15 a  $x + 2y - 16 = 0$   
 b  $y = -\frac{2}{3}x$   
 c  $C(-48, 32)$   
 d Slope of  $OA$  is  $\frac{3}{2}$ . Slope of  $OC$  is  $-\frac{2}{3}$ .  
 Lines are perpendicular.  
 e  $OA = 2\sqrt{13}$  and  $OC = 16\sqrt{13}$   
 f Area = 208  
 16 a  $d = \sqrt{50a^2} = 5a\sqrt{2}$                       b  $5\sqrt{2}$   
 c  $15\sqrt{2}$     d  $25\sqrt{2}$

- 17 a  $d = \sqrt{10x^2 - 28x + 26}$   
 b  $B(-\frac{6}{5}, -\frac{18}{5})$  and  $C(4, 12)$   
 c  $y = -\frac{1}{3}x + \frac{14}{5}$   
 d  $(\frac{7}{5}, \frac{21}{5})$   
 e 20.8  
 18 a gradient = 10.5  
 b  $C = 10.5P - 10751$   
 c When the oil production increases by 1 million tonnes, the carbon dioxide emissions increase by 10.5 million tonnes.  
 d The model is not valid for small values of  $P$ , as it is not possible to have a negative amount of carbon dioxide emissions. It is always dangerous to extrapolate beyond the range on the model in this way.

**Challenge**

- 1 130  
 2  $(\frac{78}{19}, \frac{140}{19})$   
 3  $(a, \frac{a(c-a)}{b})$

**CHAPTER 6**

**Prior knowledge check**

- 1 a  $(x+5)^2 + 3$                               b  $(x-3)^2 - 8$   
 c  $(x-6)^2 - 36$                               d  $(x + \frac{7}{2})^2 - \frac{49}{4}$   
 2 a  $y = \frac{9}{4}x - 6$                                       b  $y = -\frac{1}{2}x - \frac{3}{2}$   
 c  $y = \frac{4}{3}x + \frac{10}{3}$   
 3 a  $b^2 - 4ac = -7$  No real solutions  
 b  $b^2 - 4ac = 193$  Two real solutions  
 c  $b^2 - 4ac = 0$  One real solution  
 4  $y = -\frac{5}{6}x - \frac{3}{2}$

**Exercise 6A**

- 1 a (5, 5)    b (6, 4)    c (-1, 4)    d (0, 0)  
 e (2, 1)    f  $(-8, \frac{3}{2})$     g (4a, 0)    h  $(-\frac{u}{2}, -v)$   
 i  $(2a, a-b)$     j  $(3\sqrt{2}, 4)$     k  $(2\sqrt{2}, \sqrt{2} + 3\sqrt{3})$   
 2  $a = 10, b = 1$   
 3  $(\frac{3}{2}, 7)$   
 4  $(\frac{3a}{5}, \frac{b}{4})$   
 5 a  $(\frac{3}{2}, 3)$  or  $(1.5, 3)$     b  $y = 2x, 3 = 2 \times 1.5$   
 6 a  $(\frac{1}{8}, \frac{5}{3})$     b  $\frac{2}{3}$   
 7 Centre is  $(3, -\frac{7}{2})$ .  $3 - 2(-\frac{7}{2}) - 10 = 0$   
 8 (10, 5)  
 9  $(-7a, 17a)$





- 10  $p = 8, q = 7$   
 11  $a = -2, b = 4$

**Challenge**

- a  $p = 9, q = -1$   
 b  $y = -x + 13$   
 c AC:  $y = -x + 8$ . Lines have the same slope, so they are parallel.

**Exercise 6B**

- 1 a  $y = 2x + 3$     b  $y = -\frac{1}{3}x + \frac{47}{3}$     c  $y = \frac{5}{2}x - 25$   
 d  $y = 3$     e  $y = -\frac{3}{4}x + \frac{37}{8}$     f  $x = 9$   
 2  $y = -x + 7$   
 3  $2x - y - 8 = 0$   
 4 a  $y = -\frac{5}{3}x - \frac{13}{3}$     b  $y = 3x - 8$     c  $(\frac{11}{14}, \frac{79}{14})$   
 5  $q = -\frac{5}{4}, b = \frac{189}{8}$

**Challenge**

- a  $PR: y = -\frac{5}{2}x + \frac{9}{4}$   
 $PQ: y = -\frac{1}{4}x + \frac{33}{8}$   
 $RQ: y = 2x + 6$   
 b  $(-\frac{5}{6}, \frac{13}{3})$

**Exercise 6C**

- 1 a  $(x - 3)^2 + (y - 2)^2 = 16$   
 b  $(x + 4)^2 + (y - 5)^2 = 36$   
 c  $(x - 5)^2 + (y + 6)^2 = 12$   
 d  $(x - 2a)^2 + (y - 7a)^2 = 25a^2$   
 e  $(x + 2\sqrt{2})^2 + (y + 3\sqrt{2})^2 = 1$   
 2 a  $(-5, 4), 9$     b  $(7, 1), 4$   
 c  $(-4, 0), 5$     d  $(-4a, -a), 12a$   
 e  $(3\sqrt{5}, -\sqrt{5}), 3\sqrt{3}$   
 3 a  $(4 - 2)^2 + (8 - 5)^2 = 4 + 9 = 13$   
 b  $(0 + 7)^2 + (-2 - 2)^2 = 49 + 16 = 65$   
 c  $7^2 + (-24)^2 = 49 + 576 = 625 = 25^2$   
 d  $(6a - 2a)^2 + (-3a + 5a)^2 = 16a^2 + 4a^2 = 20a^2$   
 e  $(\sqrt{5} - 3\sqrt{5})^2 + (-\sqrt{5} - \sqrt{5})^2 = (-2\sqrt{5})^2 + (-2\sqrt{5})^2 = 20 + 20 = 40 = (2\sqrt{10})^2$   
 4  $(x - 8)^2 + (y - 1)^2 = 25$   
 5  $(x - \frac{3}{2})^2 + (y - 4)^2 = \frac{65}{4}$   
 6  $\sqrt{5}$   
 7 a  $r = 2$   
 b Distance  $PQ = PR = RQ = 2\sqrt{3}$ , three equal length sides triangle is equilateral.  
 8 a  $(x - 2)^2 + y^2 = 15$   
 b Centre  $(2, 0)$  and radius  $= \sqrt{15}$   
 9 a  $(x - 5)^2 + (y + 2)^2 = 49$   
 b Centre  $(5, -2)$  and radius  $= 7$   
 10 a Centre  $(1, -4)$ , radius  $5$   
 b Centre  $(-6, 2)$ , radius  $7$   
 c Centre  $(11, -3)$ , radius  $3\sqrt{10}$   
 d 10 Centre  $(-2.5, 1.5)$ , radius  $\frac{5\sqrt{2}}{2}$   
 e Centre  $(2, -2)$ , radius  
 11 a Centre  $(-6, -1)$   
 b  $k > -37$   
 12  $Q(-13, 28)$   
 13  $k = -2$  and  $k = 8$

**Challenge**

- 1  $k = 3, (x - 3)^2 + (y - 2)^2 = 50$   
 $k = 5, (x - 5)^2 + (y - 2)^2 = 50$   
 2  $(x + f)^2 - f^2 + (y + g)^2 - g^2 + c = 0$   
 So  $(x + f)^2 + (y + g)^2 = f^2 + g^2 - c$   
 Circle with centre  $(-f, -g)$  and radius  $\sqrt{f^2 + g^2 - c}$ .

**Exercise 6D**

- 1  $(7, 0), (-5, 0)$   
 2  $(0, 2), (0, -8)$   
 3  $(6, 10), (2, -2)$   
 4  $(4, -9), (-7, 2)$   
 5  $2x^2 - 24x + 79 = 0$  has no real solutions, therefore lines do not intersect  
 6 a  $b^2 - 4ac = 64 - 4 \times 1 \times 16 = 0$ . So there is only one point of intersection.  
 b  $(4, 7)$   
 7 a  $(0, -2), (4, 6)$     b midpoint of  $AB$  is  $(2, 2)$   
 8 a  $13$     b  $p = 1$  or  $5$   
 9 a  $A(5, 0)$  and  $B(-3, -8)$  (or vice-versa)  
 b  $y = -x - 3$   
 c  $(4, -7)$  is a solution to  $y = -x - 3$ .  
 d  $20$   
 10 a Substitute  $y = kx$  to give  
 $(k^2 + 1)x^2 - (12k + 10)x + 57 = 0$   
 $b^2 - 4ac > 0, -84k^2 + 240k - 128 > 0,$   
 $21k^2 - 60k + 32 < 0$   
 b  $0.71 < k < 2.15$   
 Exact answer is  $\frac{10}{7} - \frac{2\sqrt{57}}{21} < k < \frac{10}{7} + \frac{2\sqrt{57}}{21}$   
 11  $k < \frac{8}{17}$   
 12  $k = -20 \pm 2\sqrt{105}$

**Exercise 6E**

- 1 a  $3\sqrt{10}$   
 b Gradient of radius  $= 3x$ , gradient of line  $= -\frac{1}{3}$ , gradients are negative reciprocals and therefore perpendicular.  
 2 a  $(x - 4)^2 + (y - 6)^2 = 73$     b  $3x + 8y + 13 = 0$   
 3 a  $y = -2x - 1$   
 b Centre of circle  $(1, -3)$  satisfies  $y = -2x - 1$ .  
 4 a  $y = \frac{1}{2}x - 3$   
 b Centre of circle  $(2, -2)$  satisfies  $y = \frac{1}{2}x - 3$   
 5 a  $(-7, -6)$  satisfies  $x^2 + 18x + y^2 - 2y + 29 = 0$   
 b  $y = \frac{2}{7}x - 4$     c  $R(0, -4)$     d  $\frac{53}{2}$   
 6 a  $(0, -17), (17, 0)$   
 b  $144.5$   
 7  $y = 2x + 27$  and  $y = 2x - 13$   
 8 a  $p = 4, p = -6$   
 b  $(3, 4)$  and  $(3, -6)$   
 9 a  $(x - 11)^2 + (y + 5)^2 = 100$   
 b  $y = \frac{3}{4}x - \frac{3}{4}$   
 c  $A(8 - 4\sqrt{3}, -1 - 3\sqrt{3})$  and  $B(8 + 4\sqrt{3}, -1 + 3\sqrt{3})$   
 d  $10\sqrt{3}$   
 10 a  $y = 4x - 22$   
 b  $\alpha = 5$   
 c  $(x - 5)^2 + (y + 2)^2 = 34$   
 d  $A(5 + \sqrt{2}, -2 + 4\sqrt{2})$  and  $B(5 - \sqrt{2}, -2 - 4\sqrt{2})$

- 11 a  $P(-2, 5)$  and  $Q(4, 7)$   
 b  $y = 2x + 9$  and  $y = -\frac{1}{2}x + 9$   
 c  $y = -3x + 9$   
 d  $(0, 9)$

**Challenge**

- 1  $y = \frac{1}{2}x - 2$   
 2 a  $\angle CPR = \angle CQR = 90^\circ$  (Angle between tangent and radius)  
 $CP = CQ = \sqrt{10}$  (Radii of circle)  
 $CR = \sqrt{(6-2)^2 + (-1-1)^2} = \sqrt{20}$   
 So using Pythagoras' Theorem,  
 $PR = QR = \sqrt{20 - 10} = \sqrt{10}$   
 4 equal sides and two opposite right-angles, so  $CPRQ$  is a square  
 b  $y = \frac{1}{3}x - 3$  and  $y = -3x + 17$

**Exercise 6F**

- 1 a  $WV^2 = WU^2 + UV^2$   
 b  $(2, 3)$   
 c  $(x-2)^2 + (y-3)^2 = 41$   
 2 a  $AC^2 = AB^2 + BC^2$   
 b  $(x-5)^2 + (y-2)^2 = 25$   
 c 15  
 3 a i  $y = \frac{3}{2}x + \frac{21}{2}$       ii  $y = -\frac{2}{3}x + 4$   
 b  $(-3, 6)$   
 c  $(x+3)^2 + (y-6)^2 = 169$   
 4 a i  $y = \frac{1}{3}x + \frac{10}{3}$       ii  $x = -1$   
 b  $(x+1)^2 + (y-3)^2 = 125$   
 5  $(x-3)^2 + (y+4)^2 = 50$   
 6 a  $AB^2 + BC^2 = AC^2$   
 $AB^2 = 400, BC^2 = 100, AC^2 = 500$   
 b  $(x+2)^2 + (y-5)^2 = 125$   
 c  $D(8, 0)$  satisfies the equation of the circle.  
 7 a  $AB = BC = CD = DA = \sqrt{50}$   
 b 50  
 c  $(3, 6)$   
 8 a  $DE^2 = b^2 + 6b + 13$   
 $EF^2 = b^2 + 10b + 169$   
 $DF^2 = 200$   
 So  $b^2 + 6b + 13 + b^2 + 10b + 169 = 200$   
 $(b+9)(b-1) = 0$ ; as  $b > 0$ ,  $b = 1$   
 b  $(x+5)^2 + (y+4)^2 = 50$   
 9 a Centre  $(-1, 12)$  and radius = 13  
 b Use distance formula to find  $AB = 26$ . This is twice radius, so  $AB$  is the diameter. Other methods possible.  
 c  $C(-6, 0)$

**Mixed exercise**

- 1 a  $C(3, 6)$   
 b  $r = 10$   
 c  $(x-3)^2 + (y-6)^2 = 100$   
 d  $P$  satisfies the equation of the circle.  
 2  $(0-5)^2 + (0+2)^2 = 5^2 + 2^2 = 29 < 30$  therefore point is inside the circle  
 3 a Centre  $(0, -4)$  and radius = 3  
 b  $(0, -1)$  and  $(0, -7)$   
 c Students' own work. Equation  $x^2 = -7$  has no real solutions.  
 4 a  $P(8, 8), (8+1)^2 + (8-3)^2 = 9^2 + 5^2 = 81 + 25 = 106$   
 b  $\sqrt{106}$

- 5 a All points satisfy  $x^2 + y^2 = 1$ , therefore all lie on circle.  
 b  $AB = BC = CA$   
 6 a  $k = 1, k = -\frac{2}{5}$   
 b  $(x-1)^2 + (y-3)^2 = 13$   
 7 Substitute  $y = 3x - 9$  into the equation  
 $x^2 + px + y^2 + 4y = 20$   
 $x^2 + px + (3x-9)^2 + 4(3x-9) = 20$   
 $10x^2 + (p-42)x + 25 = 0$   
 Using the discriminant:  $(p-42)^2 - 1000 < 0$   
 $42 - 10\sqrt{10} < p < 42 + 10\sqrt{10}$   
 8  $(x-2)^2 + (y+4)^2 = 20$   
 9 a  $2\sqrt{29}$       b 12  
 10  $(-1, 0), (11, 0)$   
 11 The values of  $m$  and  $n$  are  $7 - \sqrt{105}$  and  $7 + \sqrt{105}$ .  
 12 a  $a = 6$  and  $b = 8$       b  $y = -\frac{4}{3}x + 8$       c 24  
 13 a  $p = 0, q = 24$       b  $(0, 49), (0, -1)$   
 14  $x + y + 10 = 0$   
 15 60  
 16  $l_1: y = -4x + 12$  and  $l_2: y = -\frac{8}{19}x + 12$   
 17 a  $y = \frac{1}{3}x + \frac{8}{3}$   
 b  $(x+2)^2 + (y-2)^2 = 50$   
 c 20  
 18 a  $P(-3, 1)$  and  $Q(9, -7)$   
 b  $y = \frac{3}{2}x + \frac{11}{2}$  and  $y = \frac{3}{2}x - \frac{41}{2}$   
 19 a  $y = -4x + 6$  and  $y = \frac{1}{4}x + 6$   
 b  $P(-4, 5)$  and  $Q(1, 2)$   
 c 17  
 20 a  $P(5, 16)$  and  $Q(13, 8)$   
 b  $l_2: y = \frac{1}{7}x + \frac{107}{7}$  and  $l_3: y = 7x - 83$   
 c  $l_4: y = x + 3$   
 d All 3 equations have solution  $x = 15, y = 18$   
 so  $R(15, 18)$   
 e  $\frac{200}{3}$   
 21 a  $(4, 0), (0, 12)$   
 b  $(2, 6)$   
 c  $(x-2)^2 + (y-6)^2 = 40$   
 22 a  $q = 4$   
 b  $(x + \frac{5}{2})^2 + (y-2)^2 = -\frac{65}{4}$   
 23 a  $RS^2 + ST^2 = RT^2$   
 b  $(x-2)^2 + (y+2)^2 = 61$   
 24  $(x-1)^2 + (y-3)^2 = 34$   
 25 a i  $y = -4x - 4$       ii  $x = -2$   
 b  $(x+2)^2 + (y-4)^2 = 34$

**Challenge**

- a  $x + y - 14 = 0$   
 b  $P(7, 7)$  and  $Q(9, 5)$   
 c 10

**CHAPTER 7**

**Prior knowledge check**

- 1 a  $15x^7$       b  $\frac{x}{3y}$   
 2 a  $(x-6)(x+4)$       b  $(3x-5)(x-4)$   
 3 a 8567      b 1652  
 4 a  $y = 1 - 3x$       b  $y = \frac{1}{2}x - 7$   
 5 a  $(x-1)^2 - 21$       b  $2(x+1)^2 + 13$

**Exercise 7A**

- 1 a  $4x^3 + 5x - 7$       b  $2x^4 + 9x^2 + x$   
 c  $-x^3 + 4x + \frac{6}{x}$       d  $7x^4 - x^2 - \frac{4}{x}$



- e  $\frac{4x^3 - 2x^2 + 3}{5} - \frac{x^3}{5} - \frac{2}{5x}$       f  $3x - 4x^2 - 1$   
 g  $\frac{7x^2}{5} - \frac{x^3}{5} - \frac{2}{5x}$       h  $2x - 3x^3 + 1$   
 i  $\frac{x^7}{2} - \frac{9x^3}{2} + 2x^2 - \frac{3}{x}$       j  $3x^8 + 2x^5 - \frac{4x^3}{3} + \frac{2}{3x}$
- 2 a  $x + 3$       b  $x + 4$       c  $x + 3$   
 d  $x + 7$       e  $x + 5$       f  $x + 4$   
 g  $\frac{x - 4}{x - 3}$       h  $\frac{x + 2}{x + 4}$       i  $\frac{x + 4}{x - 6}$   
 j  $\frac{2x + 3}{x - 5}$       k  $\frac{2x - 3}{x + 1}$       l  $\frac{x - 2}{x + 2}$   
 m  $\frac{2x + 1}{x - 2}$       n  $\frac{x + 4}{3x + 1}$       o  $\frac{2x + 1}{2x - 3}$
- 3  $a = 1, b = 4, c = -2$


**Exercise 7B**

- 1 a  $(x + 1)(x^2 + 5x + 3)$       b  $(x + 4)(x^2 + 6x + 1)$   
 c  $(x + 2)(x^2 - 3x + 7)$       d  $(x - 3)(x^2 + 4x + 5)$   
 e  $(x - 5)(x^2 - 3x - 2)$       f  $(x - 7)(x^2 + 2x + 8)$
- 2 a  $(x + 4)(6x^2 + 3x + 2)$       b  $(x + 2)(4x^2 + x - 5)$   
 c  $(x + 3)(2x^2 - 2x - 3)$       d  $(x - 6)(2x^2 - 3x - 4)$   
 e  $(x + 6)(-5x^2 + 3x + 5)$       f  $(x - 2)(-4x^2 + x - 1)$
- 3 a  $x^3 + 3x^2 - 4x + 1$       b  $4x^3 + 2x^2 - 3x - 5$   
 c  $-3x^3 + 3x^2 - 4x - 7$       d  $-5x^4 + 2x^3 + 4x^2 - 3x + 7$
- 4 a  $x^3 + 2x^2 - 5x + 4$       b  $x^3 - x^2 + 3x - 1$   
 c  $2x^3 + 5x + 2$       d  $3x^4 + 2x^3 - 5x^2 + 3x + 6$   
 e  $2x^4 - 2x^3 + 3x^2 + 4x - 7$       f  $4x^4 - 3x^3 - 2x^2 + 6x - 5$   
 g  $5x^3 + 12x^2 - 6x - 2$       h  $3x^4 + 5x^3 + 6$
- 5 a  $x^2 - 2x + 5$       b  $2x^2 - 6x + 1$   
 c  $-3x^2 - 12x + 2$
- 6 a  $x^2 + 4x + 12$       b  $2x^2 - x + 5$   
 c  $-3x^2 + 5x + 10$
- 7  $f(-2) = -8 + 8 + 10 - 10 = 0$ , so  $(x + 2)$  is a factor of  $x^3 + 2x^2 - 5x - 10$ . Divide  $x^3 + 2x^2 - 5x - 10$  by  $(x + 2)$  to give  $(x^2 - 5)$ . So  $x^3 + 2x^2 - 5x - 10 = (x + 2)(x^2 - 5)$ .
- 8 a  $-8$       b  $-7$       c  $-12$
- 9  $f(1) = 3 - 2 + 4 = 5$
- 10  $f(-1) = 3 + 8 + 10 + 3 - 25 = -1$
- 11  $(x + 4)(5x^2 - 20x + 7)$
- 12  $3x^2 + 6x + 4$
- 13  $x^2 + x + 1$
- 14  $x^3 - 2x^2 + 4x - 8$
- 15 14
- 16 a  $-200$       b  $(x + 2)(x - 7)(3x + 1)$
- 17 a i 30      ii 0      b  $x = -3, x = -4, x = 1$
- 18 a  $a = 1, b = 2, c = -3$   
 b  $f(x) = (2x - 1)(x + 3)(x - 1)$   
 c  $x = 0.5, x = -3, x = 1$
- 19 a  $a = 3, b = 2, c = 1$   
 b Quadratic has no real solutions so only  $(4x - 1)$  is a solution


**Exercise 7C**

- 1 a  $f(1) = 0$       b  $f(-3) = 0$       c  $f(4) = 0$   
 2  $(x - 1)(x + 3)(x + 4)$   
 3  $(x + 1)(x + 7)(x - 5)$   
 4  $(x - 5)(x - 4)(x + 2)$   
 5  $(x - 2)(2x - 1)(x + 4)$   
 6 a  $(x + 1)(x - 5)(x - 6)$       b  $(x - 2)(x + 1)(x + 2)$   
 c  $(x - 5)(x + 3)(x - 2)$


- 7 a i  $(x - 1)(x + 3)(2x + 1)$       ii




- b i  $(x - 3)(x - 5)(2x - 1)$       ii




- c i  $(x + 1)(x + 2)(3x - 1)$       ii



- d i  $(x + 2)(2x - 1)(3x + 1)$       ii



- e i  $(x - 2)(2x - 5)(2x + 3)$       ii



- 8 2  
 9  $-16$   
 10  $p = 3, q = 7$   
 11  $c = 2, d = 3$   
 12  $g = 3, h = -7$   
 13 a  $f(4) = 0$   
 b  $f(x) = (x - 4)(3x^2 + 6)$   
 For  $3x^2 + 6 = 0$ ,  $b^2 - 4ac = -72$  so there are no real roots. Therefore, 4 is the only real root of  $f(x) = 0$ .  
 14 a  $f(-2) = 0$       b  $(x + 2)(2x + 1)(2x - 3)$   
 c  $x = -2, x = -\frac{1}{2}$  and  $x = \frac{1}{2}$   
 15 a  $f(2) = 0$       b  $x = 0, x = 2, x = -\frac{1}{3}$  and  $x = \frac{1}{3}$

**Challenge**

- a  $f(1) = 2 - 5 - 42 - 9 + 54 = 0$   
 $f(-3) = 162 + 135 - 378 + 27 + 54 = 0$   
 b  $2x^4 - 5x^3 - 42x^2 - 9x + 54$   
 $= (x - 1)(x + 3)(x - 6)(2x + 3)$   
 $x = 1, x = -3, x = 6, x = -1.5$



**Exercise 7D**

- 1  $n^2 - n = n(n - 1)$   
If  $n$  is even,  $n - 1$  is odd and even  $\times$  odd = even  
If  $n$  is odd,  $n - 1$  is even and odd  $\times$  even = even
- 2  $\frac{x}{(1 + \sqrt{2})} \times \frac{(1 - \sqrt{2})}{(1 - \sqrt{2})} = \frac{x(1 - \sqrt{2})}{(1 - 2)} = \frac{x - x\sqrt{2}}{-1} = x\sqrt{2} - x$
- 3  $(x + \sqrt{y})(x - \sqrt{y}) = x^2 - x\sqrt{y} + x\sqrt{y} - y = x^2 - y$
- 4  $(2x - 1)(x + 6)(x - 5) = (2x - 1)(x^2 + x - 30)$   
 $= 2x^3 + x^2 - 61x + 30$
- 5 LHS =  $x^2 + bx$ , using completing the square,  
 $(x + \frac{b}{2})^2 - (\frac{b}{2})^2$
- 6  $x^2 + 2bx + c = 0$ , using completing the square  
 $(x + b)^2 + c - b^2 = 0$   
 $(x + b)^2 = b^2 - c$   
 $x + b = \pm \sqrt{b^2 - c}$   
 $x = -b \pm \sqrt{b^2 - c}$
- 7  $(x - \frac{2}{x})^3 = (x - \frac{2}{x})(x^2 - 4 + \frac{4}{x^2}) = x^3 - 6x + \frac{12}{x} - \frac{8}{x^3}$
- 8  $(x^3 - \frac{1}{x})(x^{\frac{3}{2}} + x^{-\frac{3}{2}}) = x^{\frac{3}{2}} + x^{\frac{3}{2}} - x^{\frac{1}{2}} - x^{-\frac{7}{2}} = x^{\frac{3}{2}} - x^{-\frac{7}{2}}$   
 $= x^{\frac{3}{2}}(x^4 - \frac{1}{x^4})$
- 9  $3n^2 - 4n + 10 = 3[n^2 - \frac{4}{3}n + \frac{10}{3}] = 3[(n - \frac{2}{3})^2 + \frac{10}{3} - \frac{4}{9}]$   
 $= 3(n - \frac{2}{3})^2 + \frac{26}{3}$   
The minimum value is  $\frac{26}{3}$  so  $3n^2 - 4n + 10$  is always positive.
- 10  $-n^2 - 2n - 3 = -[n^2 + 2n + 3] = -[(n + 1)^2 + 3 - 1]$   
 $= -(n + 1)^2 - 2$   
The maximum value is  $-2$  so  $-n^2 - 2n - 3$  is always negative.
- 11  $x^2 + 8x + 20 = (x + 4)^2 + 4$   
The minimum value is 4 so  $x^2 + 8x + 20$  is always greater than or equal to 4.
- 12  $kx^2 + 5kx + 3 = 0, b^2 - 4ac < 0, 25k^2 - 12k < 0,$   
 $k(25k - 12) < 0, 0 < k < \frac{12}{25}$   
When  $k = 0$  there are no real roots, so  $0 \leq k < \frac{12}{25}$
- 13  $px^2 - 5x - 6 = 0, b^2 - 4ac > 0, 25 + 24p > 0, p > -\frac{25}{24}$
- 14 Gradient  $AB = -\frac{1}{2}$ , gradient  $BC = 2$ ,  
Gradient  $AB \times$  gradient  $BC = -\frac{1}{2} \times 2 = -1$ ,  
so  $AB$  and  $BC$  are perpendicular.
- 15 Gradient  $AB = 3$ , gradient  $BC = \frac{1}{4}$ , gradient  $CD = 3$ ,  
gradient  $AD = \frac{1}{4}$   
Gradient  $AB =$  gradient  $CD$  so  $AB$  and  $CD$  are parallel.  
Gradient  $BC =$  gradient  $AD$  so  $BC$  and  $AD$  are parallel.
- 16 Gradient  $AB = \frac{1}{3}$ , gradient  $BC = 3$ , gradient  $CD = \frac{1}{3}$ ,  
gradient  $AD = 3$   
Gradient  $AB =$  gradient  $CD$  so  $AB$  and  $CD$  are parallel.  
Gradient  $BC =$  gradient  $AD$  so  $BC$  and  $AD$  are parallel.  
Length  $AB = \sqrt{10}, BC = \sqrt{10}, CD = \sqrt{10}$  and  $AD = \sqrt{10}$ ,  
all four sides are equal
- 17 Gradient  $AB = -3$ , gradient  $BC = \frac{1}{3}$ ,  
Gradient  $AB \times$  gradient  $BC = -3 \times \frac{1}{3} = -1$ , so  $AB$  and  $BC$   
are perpendicular  
Length  $AB = \sqrt{40}, BC = \sqrt{40}, AB = BC$
- 18  $(x - 1)^2 + y^2 = k, y = ax, (x - 1)^2 + a^2x^2 = k,$   
 $x^2(1 + a^2) - 2x + 1 - k = 0$   
 $b^2 - 4ac > 0, k > \frac{a^2}{1 + a^2}$

- 19  $x = 2$ . There is only one solution so the line  
 $4y - 3x + 26 = 0$  only touches the circle in one place so  
is the tangent to the circle.
- 20 Area of square =  $(a + b)^2 = a^2 + 2ab + b^2$   
Shaded area =  $4(\frac{1}{2}ab)$   
Area of smaller square:  $a^2 + 2ab + b^2 - 2ab$   
 $= a^2 + b^2 = c^2$

**Challenge**

- 1 The equation of the circle is  $(x - 3)^2 + (y - 5)^2 = 25$  and  
all four points satisfy this equation.
- 2  $(\frac{1}{2}(p + 1))^2 - (\frac{1}{2}(p - 1))^2 = \frac{1}{4}(p + 1)^2 - \frac{1}{4}(p - 1)^2 = \frac{1}{4}(4p) = p$

**Exercise 7E**

- 1 3, 4, 5, 6, 7 and 8 are not divisible by 10
- 2 3, 5, 7, 11, 13, 17, 19, 23 are prime numbers. 9, 15, 21,  
25, are the product of two prime numbers.
- 3  $2^2 + 3^2 =$  odd,  $3^2 + 4^2 =$  odd,  $4^2 + 5^2 =$  odd,  $5^2 + 6^2 =$  odd,  
 $6^2 + 7^2 =$  odd
- 4  $(3n)^2 = 27n^2 = 9n(3n^2)$  which is a multiple of 9  
 $(3n + 1)^2 = 27n^2 + 27n + 9n + 1 = 9n(3n^2 + 3n + 1) + 1$   
which is one more than a multiple of 9  
 $(3n + 2)^2 = 27n^2 + 54n^2 + 36n + 8 = 9n(3n^2 + 6n + 4) + 8$   
which is one less than a multiple of 9
- 5 a For example, when  $n = 2, 2^4 - 2 = 14, 14$  is not  
divisible by 4.  
b Any square number  
c For example, when  $n = \frac{1}{2}$   
d For example, when  $n = 1$
- 6 a Assuming that  $x$  and  $y$  are positive  
b e.g.  $x = 0, y = 0$
- 7  $(x + 5)^2 \geq 0$  for all real values of  $x$ , and  
 $(x + 5)^2 + 2x + 11 = (x + 6)^2$ , so  $(x + 6)^2 \geq 2x + 11$
- 8 If  $a^2 + 1 \geq 2a$  ( $a$  is positive, so multiplying both sides  
by  $a$  does not reverse the inequality), then  
 $a^2 - 2a + 1 \geq 0$ , and  $(a - 1)^2 \geq 0$ , which we know is  
true.
- 9 a  $(p + q)^2 = p^2 + 2pq + q^2 = (p + q)^2 + 4pq$   
 $(p - q)^2 \geq 0$  since it is a square, so  $(p + q)^2 \geq 4pq$   
 $p > 0, q > 0 \Rightarrow p + q > 0 \Rightarrow p + q \geq \sqrt{4pq}$   
b e.g.  $p = q = -1: p + q = -2, \sqrt{4pq} = 2$
- 10 a Starts by assuming the inequality is true:  
i.e. negative  $\geq$  positive  
b e.g.  $x = y = -1: x + y = -2, \sqrt{x^2 + y^2} = \sqrt{2}$   
c  $(x + y)^2 = x^2 + 2xy + y^2 \geq x^2 + y^2$  since  $x > 0,$   
 $y > 0 \Rightarrow 2xy > 0$   
As  $x + y > 0$ , can take square roots:  $x + y \geq \sqrt{x^2 + y^2}$

**Mixed exercise**

- 1 a  $x^3 - 7$       b  $\frac{x + 4}{x - 1}$       c  $\frac{2x - 1}{2x + 1}$
- 2  $3x^2 + 5$
- 3  $2x^2 - 2x + 5$
- 4 a When  $x = 3, 2x^3 - 2x^2 - 17x + 15 = 0$   
b  $A = 2, B = 4, C = -5$
- 5 a When  $x = 2, x^3 + 4x^2 - 3x - 18 = 0$   
b  $p = 1, q = 3$
- 6  $(x - 2)(x + 4)(2x - 1)$
- 7 7
- 8 a  $p = 1, q = -15$       b  $(x + 3)(2x - 5)$
- 9 a  $r = 3, s = 0$       b  $x(x + 1)(x + 3)$
- 10 a  $(x - 1)(x + 5)(2x + 1)$       b  $-5, -\frac{1}{2}, 1$





- 11 a When  $x = 2$ ,  $x^3 + x^2 - 5x - 2 = 0$   
 b  $2, -\frac{3}{2} \pm \frac{\sqrt{5}}{2}$
- 12  $\frac{1}{2}, 3$
- 13 a When  $x = -4$ ,  $f(x) = 0$   
 b  $(x + 4)(x - 5)(x - 1)$
- 14 a  $f(\frac{2}{3}) = 0$ , therefore  $(3x - 2)$  is a factor of  $f(x)$   
 $a = 2, b = 7$  and  $c = 3$   
 b  $(3x - 2)(2x + 1)(x + 3)$   
 c  $x = \frac{2}{3}, -\frac{1}{2}, -3$
- 15  $\frac{x - y}{(\sqrt{x} - \sqrt{y})} \times \frac{(\sqrt{x} + \sqrt{y})}{(\sqrt{x} + \sqrt{y})} = \frac{x\sqrt{x} + x\sqrt{y} - y\sqrt{x} - y\sqrt{y}}{x - y} = \sqrt{x} + \sqrt{y}$
- 16  $n^2 - 8n + 20 = (n - 4)^2 + 4$ , 4 is the minimum value so  $n^2 - 8n + 20$  is always positive
- 17 Gradient  $AB = \frac{1}{2}$ , gradient  $BC = -2$ , gradient  $CD = \frac{1}{2}$ , gradient  $AD = -2$   
 $AB$  and  $BC$ ,  $BC$  and  $CD$ ,  $CD$  and  $AD$  and  $AB$  and  $AD$  are all perpendicular  
 Length  $AB = \sqrt{5}$ ,  $BC = \sqrt{5}$ ,  $CD = \sqrt{5}$  and  $AD = \sqrt{5}$ , all four sides are equal
- 18  $1 + 3 = \text{even}$ ,  $3 + 5 = \text{even}$ ,  $5 + 7 = \text{even}$ ,  $7 + 9 = \text{even}$
- 19 For example when  $n = 6$
- 20  $(x - \frac{1}{x})(x^{\frac{1}{2}} + x^{\frac{3}{2}}) = x^{\frac{3}{2}} + x^{\frac{5}{2}} - x^{\frac{1}{2}} - x^{\frac{3}{2}} = x^{\frac{1}{2}}(x^2 - \frac{1}{x^2})$
- 21 RHS =  $(x + 4)(x - 5)(2x + 3) = (x + 4)(2x^2 - 7x - 15) = 2x^3 + x^2 - 43x - 60 = \text{LHS}$
- 22  $x^2 - kx + k = 0$ ,  $b^2 - 4ac = 0$ ,  $k^2 - 4k = 0$ ,  $k(k - 4) = 0$ ,  $k = 4$ .
- 23 The distance between opposite edges  
 $= 2\left(\left(\sqrt{3}\right)^2 - \left(\frac{\sqrt{3}}{2}\right)^2\right) = 2\left(3 - \frac{3}{4}\right) = \frac{9}{2}$  which is rational.
- 24 a  $(2n + 2)^2 - (2n)^2 = 8n + 4 = 4(2n + 1)$  is always divisible by 4.  
 b Yes,  $(2n + 1)^2 - (2n - 1)^2 = 8n$  which is always divisible by 4.
- 25 a The assumption is that  $x$  is positive  
 b  $x = 0$

**Challenge**

- 1 a Perimeter of inside square =  $4\left(\sqrt{\left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^2}\right) = \frac{4}{\sqrt{2}} = 2\sqrt{2}$   
 Perimeter of outside square = 4,  
 therefore  $2\sqrt{2} < \pi < 4$ .  
 b Perimeter of inside hexagon = 3  
 Perimeter of outside hexagon =  $6 \times \frac{\sqrt{3}}{3} = 2\sqrt{3}$ ,  
 therefore  $3 < \pi < 2\sqrt{3}$
- 2  $ax^3 + bx^2 + cx + d \div (x - p) = ax^2 + (b + ap)x + (c + bp + ap^2)$  with remainder  $d + cp + bp^2 + ap^3$   
 $f(p) = ap^3 + bp^2 + cp + d = 0$ , which matches the remainder, so  $(x - p)$  is a factor of  $f(x)$ .

**CHAPTER 8**

**Prior knowledge check**

- 1 a  $4x^2 - 12xy - 9y^2$       b  $x^3 + 3x^2y + 3xy^2 + y^3$   
 c  $8 + 12x + 6x^2 + x^3$
- 2 a  $-8x^3$       b  $\frac{1}{81x^4}$       c  $\frac{4}{25}x^2$       d  $\frac{27}{x^3}$
- 3 a  $5\sqrt{x}$       b  $\frac{1}{16\sqrt{x^2}}$       c  $\frac{10}{3\sqrt{x}}$       d  $\frac{16\sqrt{x^4}}{81}$

**Exercise 8A**

- 1 a 4th row      b 16th row  
 c  $(n + 1)$ th row      d  $(n + 5)$ th row
- 2 a  $x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$   
 b  $p^5 + 5p^4q + 10p^3q^2 + 10p^2q^3 + 5pq^4 + q^5$   
 c  $a^3 - 3a^2b + 3ab^2 - b^3$   
 d  $x^3 + 12x^2 + 48x + 64$   
 e  $16x^4 - 96x^3 + 216x^2 - 216x + 81$   
 f  $a^5 + 10a^4 + 40a^3 + 80a^2 + 80a + 32$   
 g  $81x^4 - 432x^3 + 864x^2 - 768x + 256$   
 h  $16x^4 - 96x^3y + 216x^2y^2 - 216xy^3 + 81y^4$
- 3 a 16      b -10      c 8      d 1280  
 e 160      f -2      g 40      h -96
- 4  $1 + 9x + 30x^2 + 44x^3 + 24x^4$   
 5  $8 + 12y + 6y^2 + y^3$ ,  $8 + 12x - 6x^2 - 11x^3 + 3x^4 + 3x^5 - x^6$   
 6  $\pm 3$   
 7  $\frac{5}{2}, -1$   
 8  $12p$   
 9  $500 + 25X + \frac{X^2}{2}$

**Challenge**

$\frac{3}{4}$

**Exercise 8B**

- 1 a 24      b 362 880      c 720      d 210  
 2 a 6      b 15      c 20      d 5  
 e 45      f 126
- 3 a 5005      b 120      c 184 756      d 1140  
 e 2002      f 8568
- 4  $a = {}^4C_1, b = {}^5C_2, c = {}^6C_2, d = {}^6C_3$   
 5 330  
 6 a 120, 210      b 960  
 7 a 286, 715      b 57 915  
 8 0.1762 to 4 decimal places. Whilst it seems a low probability, there is more chance of the coin landing on 10 heads than any other amount of heads.
- 9 a  ${}^nC_1 = \frac{n!}{1!(n-1)!} = \frac{1 \times 2 \times \dots \times (n-2) \times (n-1) \times n}{1 \times 1 \times 2 \times \dots \times (n-3) \times (n-2) \times (n-1)} = n$   
 b  ${}^nC_2 = \frac{n!}{2!(n-2)!} = \frac{1 \times 2 \times \dots \times (n-2) \times (n-1) \times n}{1 \times 2 \times 1 \times 2 \times \dots \times (n-3) \times (n-2)} = \frac{n(n-1)}{2}$
- 10  $a = 37$   
 11  $p = 17$

**Challenge**

- a  ${}^{10}C_3 = \frac{10!}{3!7!} = 120$  and  ${}^{10}C_7 = \frac{10!}{7!3!} = 120$   
 b  ${}^{14}C_5 = \frac{14!}{5!9!} = 2002$  and  ${}^{14}C_9 = \frac{14!}{9!5!} = 2002$   
 c The two answers for part a are the same and the two answers for part b are the same.  
 d  ${}^nC_r = \frac{n!}{r!(n-r)!}$  and  ${}^nC_{n-r} = \frac{n!}{(n-r)!r!}$ , therefore  ${}^nC_r = {}^nC_{n-r}$

**Exercise 8C**

- 1 a  $1 + 4x + 6x^2 + 4x^3 + x^4$   
 b  $81 + 108x + 54x^2 + 12x^3 + x^4$   
 c  $256 - 256x + 96x^2 - 16x^3 + x^4$

- d  $x^6 + 12x^5 + 60x^4 + 160x^3 + 240x^2 + 192x + 64$   
 e  $1 + 8x + 24x^2 + 32x^3 + 16x^4$   
 f  $1 - 2x + \frac{3}{2}x^2 - \frac{1}{2}x^3 + \frac{1}{16}x^4$
- 2 a  $1 + 10x + 45x^2 + 120x^3$   
 b  $1 - 10x + 40x^2 - 80x^3$   
 c  $1 + 18x + 135x^2 + 540x^3$   
 d  $256 - 1024x + 1792x^2 - 1792x^3$   
 e  $1024 - 2560x + 2880x^2 - 1920x^3$   
 f  $2187 - 5103x + 5103x^2 - 2835x^3$
- 3 a  $64x^6 + 192x^5y + 240x^4y^2 + 160x^3y^3$   
 b  $32x^5 + 240x^4y + 720x^3y^2 + 1080x^2y^3$   
 c  $p^8 - 8p^7q + 28p^6q^2 - 56p^5q^3$   
 d  $729x^6 - 1458x^5y + 1215x^4y^2 - 540x^3y^3$   
 e  $x^8 + 16x^7y + 112x^6y^2 + 448x^5y^3$   
 f  $512x^9 - 6912x^8y + 41472x^7y^2 - 145152x^6y^3$
- 4 a  $1 + 8x + 28x^2 + 56x^3$   
 b  $1 - 12x + 60x^2 - 160x^3$   
 c  $1 + 5x + \frac{45}{4}x^2 + 15x^3$   
 d  $1 - 15x + 90x^2 - 270x^3$   
 e  $128 + 448x + 672x^2 + 560x^3$   
 f  $27 - 54x + 36x^2 - 8x^3$   
 g  $64 - 576x + 2160x^2 - 4320x^3$   
 h  $256 + 256x + 96x^2 + 16x^3$   
 i  $128 + 2240x + 16800x^2 + 70000x^3$
- 5  $64 - 192x + 240x^2$   
 6  $243 - 810x + 1080x^2$   
 7  $x^5 + 5x^3 + 10x + \frac{10}{x} + \frac{5}{x^3} + \frac{1}{x^5}$

Challenge

- a  $(a + b)^4 = a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4$   
 $(a - b)^4 = a^4 - 4a^3b + 6a^2b^2 - 4ab^3 + b^4$   
 $(a + b)^4 - (a - b)^4 = 8a^3b + 8ab^3 = 8ab(a^2 + b^2)$   
 b  $82896 = 2^4 \times 3 \times 11 \times 157$

Exercise 8D

- 1 a 90                      b 80                      c -20  
 d 1080                    e 120                    f -4320  
 g 1140                    h -241920              i -2.5  
 j 354.375                k -224                   l 3.90625
- 2  $a = \pm \frac{1}{2}$   
 3  $b = -2$   
 4  $1, \frac{5 \pm \sqrt{105}}{8}$
- 5 a  $p = 5$                 b -10                    c -80  
 6 a  $5^{30} + 5^{29} \times 30px + 5^{28} \times 435p^2x^2$   
 b  $p = 10$   
 7 a  $1 + 10qx + 45q^2x^2 + 120q^3x^3$   
 b  $q = \pm 3$   
 8 a  $1 + 11px + 55p^2x^2$   
 b  $p = 7, q = 2695$   
 9 a  $1 + 15px + 105p^2x^2$   
 b  $p = -\frac{5}{7}, q = 10\frac{5}{7}$   
 10  $\frac{q}{p} = 2.1$

Challenge

- a 314928                b 43750

Exercise 8E

- 1 a  $1 - 0.6x + 0.15x^2 - 0.02x^3$   
 b 0.94148

- 2 a  $1024 + 1024x + 460.8x^2 + 122.88x^3$   
 b 1666.56
- 3  $(1 - 3x)^5 = 1^5 + \binom{5}{1}1^4(-3x)^1 + \binom{5}{2}1^3(-3x)^2 = 1 - 15x + 90x^2$   
 $(2 + x)(1 - 3x)^5 = (2 + x)(1 - 15x + 90x^2)$   
 $= 2 - 30x + 180x^2 + x - 15x^2 + 90x^3 \approx 2 - 29x + 165x^2$   
 4  $a = 162, b = 135, c = 0$   
 5 a  $1 + 16x + 112x^2 + 448x^3$   
 b  $x = 0.01, 1.02^8 \approx 1.171648$   
 6 a  $1 - 150x + 10875x^2 - 507500x^3$   
 b 0.860368  
 c 0.860384, 0.0019%  
 7 a  $59049 - 39366x + 11809.8x^2$   
 b Substitute  $x = 0.1$  into the expansion.  
 8 a  $1 - 15x + 90x^2 - 270x^3$   
 b  $(1 + x)(1 - 3x)^5 \approx (1 + x)(1 - 15x) \approx 1 - 14x$   
 9 a So that higher powers of  $p$  can be ignored as they tend to 0  
 b  $1 + 200p - 19900p^2$   
 c  $p = 0.000417$  (3 s.f.)

Mixed exercise

- 1 a 455, 1365            b 3640  
 2  $a = 28$   
 3 a 0.0148              b 0.0000000000349            c 0.166  
 4 a  $p = 16$               b 270                      c -1890  
 5  $A = 8192, B = -53248, C = 159744$   
 6 a  $1 - 20x + 180x^2 - 960x^3$   
 b 0.81704,  $x = 0.01$   
 7 a  $1024 - 15360x + 103680x^2 - 414720x^3$   
 b 880.35  
 8 a  $81 + 216x + 216x^2 + 96x^3 + 16x^4$   
 b  $81 - 216x + 216x^2 - 96x^3 + 16x^4$   
 c 1154  
 9 a  $n = 8$                 b  $\frac{35}{8}$   
 10 a  $81 + 1080x + 5400x^2 + 12000x^3 + 10000x^4$   
 b 1012054108081,  $x = 100$   
 11 a  $1 + 24x + 264x^2 + 1760x^3$             b 1.26816  
 c 1.268241795                                  d 0.00645% (3 sf)
- 12  $x^5 - 5x^3 + 10x - \frac{10}{x} + \frac{5}{x^3} - \frac{1}{x^5}$
- 13 a  $\binom{n}{2}(2k)^{n-2} = \binom{n}{3}(2k)^{n-3}$   
 $\frac{n!(2k)^{n-2}}{2!(n-2)!} = \frac{n!(2k)^{n-3}}{3!(n-3)!}$   
 $\frac{2k}{n-2} = \frac{1}{3}$   
 So  $n = 6k + 2$   
 b  $\frac{4096}{729} + \frac{2048}{81}x + \frac{1280}{27}x^2 + \frac{1280}{27}x^3$
- 14 a  $64 + 192x + 240x^2 + 160x^3 + 60x^4 + 12x^5 + x^6$   
 b  $k = 1560$   
 15 a  $k = 1.25$             b 3500  
 16 a  $A = 64, B = 160, C = 20$             b  $x = \pm\sqrt{\frac{3}{2}}$   
 17 a  $p = 1.5$             b 50.625  
 18 672  
 19 a  $128 + 448px + 672p^2x^2$   
 b  $p = 5, q = 16800$   
 20 a  $1 - 12px + 66p^2x^2$   
 b  $p = -1\frac{1}{11}, q = 13\frac{1}{11}$



- 21 a  $128 + 224x + 168x^2$   
 b Substitute  $x = 0.1$  into the expansion.  
 22  $k = \frac{1}{2}$

**Challenge**

- 1  $540 - 405p = 0, p = \frac{4}{3}$   
 2  $-4704$

**CHAPTER 9**

**Prior knowledge check**

- 1 a 3.10 cm      b 9.05 cm  
 2 a  $25.8^\circ$       b  $77.2^\circ$   
 3 a graph of  $x^2 + 3x$       b graph of  $(x + 2)^2 + 3(x + 2)$   
 c graph of  $x^2 + 3x - 3$       b graph of  $(0.5x)^2 + 3(0.5x)$

**Exercise 9A**

- 1 a 3.19 cm      b 1.73 cm ( $\sqrt{3}$  cm)      c 9.85 cm  
 d 4.31 cm      e 6.84 cm      f 9.80 cm  
 2 a  $108(2)^\circ$       b  $90^\circ$       c  $60^\circ$   
 d  $52.6^\circ$       e  $137^\circ$       f  $72.2^\circ$   
 3 192 km  
 4 11.2 km  
 5  $128.5^\circ$  or  $031.5^\circ$  (Angle  $BAC = 48.5^\circ$ )  
 6 302 yards (301.5...)  
 7 Using the cosine rule  $\frac{5^2 + 4^2 - 6^2}{2 \times 5 \times 4} = \frac{1}{8}$   
 8 Using the cosine rule  $\frac{2^2 + 3^2 - 4^2}{2 \times 2 \times 3} = -\frac{1}{4}$   
 9  $ACB = 22.3^\circ$   
 10  $ABC = 108(4)^\circ$   
 11  $104^\circ$  ( $104.48^\circ$ )  
 12 4.4 cm  
 13 42 cm  
 14 a  $y^2 = (5 - x)^2 + (4 + x)^2 - 2(5 - x)(4 + x) \cos 120^\circ$   
 $= 25 - 10x + x^2 + 16 + 8x + x^2 - 2(20 + x - x^2) \left(-\frac{1}{2}\right)$   
 $= x^2 - x + 61$   
 b Minimum  $AC^2 = 60.75$ ; it occurs for  $x = \frac{1}{2}$   
 15 a  $\cos \angle ABC = \frac{x^2 + 5^2 - (10 - x)^2}{2x \times 5}$   
 $= \frac{20x - 75}{10x} = \frac{4x - 15}{2x}$   
 b 3.5  
 16  $65.3^\circ$   
 17 a 28.7 km      b  $056.6^\circ$

**Exercise 9B**

- 1 a 15.2 cm      b 9.57 cm      c 8.97 cm      d 4.61 cm  
 2 a  $x = 84^\circ, y = 6.32$   
 b  $x = 13.5, y = 16.6$   
 c  $x = 85^\circ, y = 13.9$   
 d  $x = 80^\circ, y = 6.22$  (isosceles triangle)  
 e  $x = 6.27, y = 7.16$   
 f  $x = 4.49, y = 7.49$  (right-angled)  
 3 a  $36.4^\circ$       b  $35.8^\circ$       c  $40.5^\circ$       d  $130^\circ$   
 4 a  $48.1^\circ$       b  $45.6^\circ$       c  $14.8^\circ$       d  $48.7^\circ$   
 e  $86.5^\circ$       f  $77.4^\circ$   
 5 a 1.41 cm ( $\sqrt{2}$  cm)      b 1.93 cm  
 6  $QPR = 50.6^\circ, PQR = 54.4^\circ$   
 7 a  $x = 43.2^\circ, y = 5.02$  cm      b  $x = 101^\circ, y = 15.0$  cm  
 c  $x = 6.58$  cm,  $y = 32.1^\circ$       d  $x = 54.6^\circ, y = 10.3$  cm  
 e  $x = 21.8^\circ, y = 3.01$       f  $x = 45.9^\circ, y = 3.87^\circ$   
 8 a 6.52 km      b 3.80 km  
 9 a 7.31 cm      b 1.97 cm


- 10 a  $66.3^\circ$       b 148 m

- 11 Using the sine rule,  $x = \frac{4\sqrt{2}}{2 + \sqrt{2}}$ ; rationalising  
 $x = \frac{4\sqrt{2}(2 - \sqrt{2})}{2} = 4\sqrt{2} - 4 = 4(\sqrt{2} - 1)$ .

- 12 a 36.5 m  
 b That the angles have been measured from ground level

**Exercise 9C**

- 1 a  $70.5^\circ, 109^\circ$  ( $109.5^\circ$ )  
 b



- 2 a  $x = 74.6^\circ, y = 65.4^\circ$   
 $x = 105^\circ, y = 34.6^\circ$   
 b  $x = 59.8^\circ, y = 48.4$  cm  
 $x = 120^\circ, y = 27.3$  cm  
 c  $x = 56.8^\circ, y = 4.37$  cm  
 $x = 23.2^\circ, y = 2.06$  cm  
 3 a 5 cm ( $ACB = 90^\circ$ )      b  $24.6^\circ$   
 c  $45.6^\circ, 134(4)^\circ$   
 4 2.96 cm  
 5 In one triangle  $ABC = 101^\circ$  ( $100.9^\circ$ ); in the other  $BAC = 131^\circ$  ( $130.9^\circ$ )  
 6 a  $62.0^\circ$       b The swing is symmetrical

**Exercise 9D**

- 1 a 23.7 cm<sup>2</sup>      b 4.31 cm<sup>2</sup>      c 20.2 cm<sup>2</sup>  
 2 a  $x = 41.8^\circ$  or  $138(2)^\circ$   
 b  $x = 26.7^\circ$  or  $153(3)^\circ$   
 c  $x = 60^\circ$  or  $120^\circ$   
 3 275(3) m (third side = 135.3 m)  
 4 3.58  
 5 a Area =  $\frac{1}{2}(x + 2)(5 - x) \sin 30^\circ$   
 $= \frac{1}{2}(10 + 3x - x^2) \times \frac{1}{2}$   
 $= \frac{1}{4}(10 + 3x - x^2)$   
 b Maximum  $A = 3\frac{1}{16}$ , when  $x = 1\frac{1}{2}$   
 6 a  $\frac{1}{2}x(5 + x) \sin 150^\circ = \frac{15}{4}$   
 $\frac{1}{2}(5x + x^2) \times \frac{1}{2} = \frac{15}{4}$   
 $5x + x^2 = 15$   
 $x^2 + 5x - 15 = 0$   
 b 2.11




**Exercise 9E**

- 1 a  $x = 37.7^\circ, y = 86.3^\circ, z = 6.86$   
 b  $x = 48^\circ, y = 19.5, z = 14.6$   
 c  $x = 30^\circ, y = 11.5, z = 11.5$   
 d  $x = 21.0^\circ, y = 29.0^\circ, z = 8.09$   
 e  $x = 93.8^\circ, y = 56.3^\circ, z = 29.9^\circ$   
 f  $x = 97.2^\circ, y = 41.4^\circ, z = 41.4^\circ$



- g  $x = 45.3^\circ, y = 94.7^\circ, z = 14.7$   
 or  $x = 135^\circ, y = 5.27^\circ, z = 1.36$
- h  $x = 7.07, y = 73.7^\circ, z = 61.3^\circ$   
 or  $x = 7.07, y = 106^\circ, z = 28.7^\circ$
- i  $x = 49.8^\circ, y = 9.39, z = 37.0^\circ$
- 2 a  $ABC = 108^\circ, ACB = 32.4^\circ, AC = 15.1$  cm  
 Area =  $41.2$  cm<sup>2</sup>  
 b  $BAC = 41.5^\circ, ABC = 28.5^\circ, AB = 9.65$  cm  
 Area =  $15.7$  cm<sup>2</sup>
- 3 a 8 km      b  $060^\circ$
- 4 107 km
- 5 12 km
- 6 a 5.44      b 7.95      c  $36.8^\circ$
- 7 a  $AB + BC > AC \Rightarrow x + 6 > 7 \Rightarrow x > 1$ ;  
 $AC + AB > BC \Rightarrow 11 > x + 2 \Rightarrow x < 9$   
 b i  $x = 6.08$  from  $x^2 = 37$   
 Area =  $14.0$  cm<sup>2</sup>  
 ii  $x = 7.23$  from  $x^2 - 4(\sqrt{2} - 1)x - (29 + 8\sqrt{2}) = 0$   
 Area =  $13.1$  cm<sup>2</sup>
- 8 a  $x = 4$       b  $4.68$  cm<sup>2</sup>
- 9  $AC = 1.93$  cm
- 10 a  $AC^2 = (2 - x)^2 + (x + 1)^2 - 2(2 - x)(x + 1) \cos 120^\circ$   
 $= (4 - 4x + x^2) + (x^2 + 2x + 1) - 2(-x^2 + x + 2) \left(-\frac{1}{2}\right)$   
 $= x^2 - x + 7$   
 b  $\frac{1}{2}$
- 11  $4\sqrt{10}$
- 12  $AC = 1\frac{2}{3}$  cm and  $BC = 6$  cm  
 Area =  $5.05$  cm<sup>2</sup>
- 13 a  $61.3^\circ$       b  $78.9$  cm<sup>2</sup>
- 14 a  $DAB = 136.3^\circ, BCD = 50.1^\circ$   
 b  $13.1$  m<sup>2</sup>  
 c  $5.15$  m
- 15  $34.2$  cm<sup>2</sup>


**Exercise 9F**




- 4 a  $-30^\circ$   
 b i  $-120^\circ$       ii  $-60^\circ, 120^\circ$   
 c i  $135^\circ$       ii  $-45^\circ, -135^\circ$

**Exercise 9G**

- 1 a i 1,  $x = 0^\circ$       ii  $-1, x = 180^\circ$   
 b i 4,  $x = 90^\circ$       ii  $-4, x = 270^\circ$   
 c i 1,  $x = 0^\circ$       ii  $-1, x = 180^\circ$   
 d i 4,  $x = 90^\circ$       ii 2,  $x = 270^\circ$   
 e i 1,  $x = 270^\circ$       ii  $-1, x = 90^\circ$   
 f i 1,  $x = 30^\circ$       ii  $-1, x = 90^\circ$




- 3 a The graph of  $y = -\cos \theta$  is the graph of  $y = \cos \theta$  reflected in the  $\theta$ -axis



- Meets  $\theta$ -axis at  $(90^\circ, 0), (270^\circ, 0)$   
 Meets  $y$ -axis at  $(0^\circ, -1)$   
 Maximum at  $(180^\circ, 1)$   
 Minimum at  $(0^\circ, -1)$  and  $(360^\circ, -1)$

- b The graph of  $y = \frac{1}{3} \sin \theta$  is the graph of  $y = \sin \theta$  stretched by a scale factor  $\frac{1}{3}$  in the  $y$  direction.




- Meets  $\theta$ -axis at  $(0^\circ, 0), (180^\circ, 0), (360^\circ, 0)$   
 Meets  $y$ -axis at  $(0^\circ, 0)$   
 Maximum at  $(90^\circ, \frac{1}{3})$   
 Minimum at  $(270^\circ, -\frac{1}{3})$






- c The graph of  $y = \sin \frac{1}{3}\theta$  is the graph of  $y = \sin \theta$  stretched by a scale factor 3 in the  $\theta$  direction.



Only meets axis at origin  
Maximum at  $(270^\circ, 1)$


- d The graph of  $y = \tan(\theta - 45^\circ)$  is the graph of  $\tan \theta$  translated by  $45^\circ$  in the positive  $\theta$  direction.



Meets  $\theta$ -axis at  $(45^\circ, 0)$ ,  $(225^\circ, 0)$   
Meets  $y$ -axis at  $(0^\circ, -1)$


(Asymptotes at  $\theta = 135^\circ$  and  $\theta = 315^\circ$ )

- 4 a This is the graph of  $y = \sin \theta$  stretched by scale factor  $-2$  in the  $y$ -direction (i.e. reflected in the  $\theta$ -axis and scaled by 2 in the  $y$ -direction).




Meets  $\theta$ -axis at  $(-180^\circ, 0)$ ,  $(0, 0)$ ,  $(180^\circ, 0)$   
Maximum at  $(-90^\circ, 2)$   
Minimum at  $(90^\circ, -2)$ .

- b This is the graph of  $y = \tan \theta$  translated by  $180^\circ$  in the negative  $\theta$  direction.



As  $\tan \theta$  has a period of  $180^\circ$   
 $\tan(\theta + 180^\circ) = \tan \theta$   
Meets  $\theta$ -axis at  $(-180^\circ, 0)$ ,  $(0, 0)$ ,  $(180^\circ, 0)$   
Meets  $y$ -axis at  $(0, 0)$

- c This is the graph of  $y = \cos \theta$  stretched by scale factor  $\frac{1}{4}$  horizontally.




Meets  $\theta$ -axis at  $(-157\frac{1}{2}^\circ, 0)$ ,  $(-112\frac{1}{2}^\circ, 0)$ ,  $(-67\frac{1}{2}^\circ, 0)$ ,  
 $(-22\frac{1}{2}^\circ, 0)$ ,  $(22\frac{1}{2}^\circ, 0)$ ,  $(67\frac{1}{2}^\circ, 0)$ ,  $(112\frac{1}{2}^\circ, 0)$ ,  $(157\frac{1}{2}^\circ, 0)$

Meets  $y$ -axis at  $(0, 1)$

Maxima at  $(-180^\circ, 1)$ ,  $(-90^\circ, 1)$ ,  $(0, 1)$ ,  $(90^\circ, 1)$ ,  $(180^\circ, 1)$

Minima at  $(-135^\circ, -1)$ ,  $(-45^\circ, -1)$ ,  $(45^\circ, -1)$ ,  $(135^\circ, -1)$

- d This is the graph of  $y = \sin \theta$  reflected in the  $y$ -axis. (This is the same as  $y = -\sin \theta$ .)




Meets  $\theta$ -axis at  $(-180^\circ, 0)$ ,  $(0^\circ, 0)$ ,  $(180^\circ, 0)$


Maximum at  $(-90^\circ, 1)$

Minimum at  $(90^\circ, -1)$


- 5 a Period =  $720^\circ$




- b Period =  $360^\circ$




- c Period =  $180^\circ$




d Period =  $90^\circ$




6 a i  $y = \cos(-\theta)$  is a reflection of  $y = \cos \theta$  in the  $y$ -axis, which is the same curve, so  $\cos \theta = \cos(-\theta)$ .




ii  $y = \sin(-\theta)$  is a reflection of  $y = \sin \theta$  in the  $y$ -axis.



$y = -\sin(-\theta)$  is a reflection of  $y = \sin(-\theta)$  in the  $\theta$ -axis, which is the graph of  $y = \sin \theta$ , so  $-\sin(-\theta) = \sin \theta$ .



iii  $y = \sin(\theta - 90^\circ)$  is the graph of  $y = \sin \theta$  translated by  $90^\circ$  to the right, which is the graph of  $y = -\cos \theta$ , so  $\sin(\theta - 90^\circ) = -\cos \theta$ .



b  $\sin(90^\circ - \theta)$   
 $= -\sin(-90^\circ - \theta) = -\sin(\theta - 90^\circ)$   
 using (a) (ii)  
 $= -(-\cos \theta)$  using (a) (iii)  
 $= \cos \theta$

c Using (a)(i)  $\cos(90^\circ - \theta) = \cos(-90^\circ - \theta)$   
 $= \cos(\theta - 90^\circ)$ , but  $\cos(\theta - 90^\circ) = \sin \theta$ ,  
 so  $\cos(90^\circ - \theta) = \sin \theta$


7 a  $(-300^\circ, 0)$ ,  $(-120^\circ, 0)$ ,  $(60^\circ, 0)$ ,  $(240^\circ, 0)$

b  $(0^\circ, \frac{\sqrt{3}}{2})$

8 a  $y = \sin(x + 60^\circ)$

b Yes – could also be a translation of the cos graph, e.g.  $y = \cos(x - 30^\circ)$

9 a



b Between 1 pm and 5 pm

**Mixed exercise**

1 a  $155^\circ$  b  $13.7$  cm

2 a  $x = 49.5^\circ$ , area =  $1.37$  cm<sup>2</sup>

b  $x = 55.2^\circ$ , area =  $10.6$  cm<sup>2</sup>

c  $x = 117^\circ$ , area =  $6.66$  cm<sup>2</sup>

3  $6.50$  cm<sup>2</sup>

4 a  $36.1$  cm<sup>2</sup> b  $12.0$  cm<sup>2</sup>

5 a  $5$  b  $\frac{25\sqrt{3}}{2}$  cm<sup>2</sup>

6 area =  $\frac{1}{2}ab \sin C$

$$1 = \frac{1}{2} \times 2\sqrt{2} \sin C$$

$$\frac{1}{\sqrt{2}} = \sin C \Rightarrow C = 45^\circ$$

Use the cosine rule to find the other side:

$$x^2 - 2^2 + (\sqrt{2})^2 - 2 \times 2\sqrt{2} \cos C \Rightarrow x = \sqrt{2}$$

So the triangle is isosceles, with two  $45^\circ$  angles, thus is also right-angled.

7 a  $AC = \sqrt{5}$ ,  $AB = \sqrt{18}$ ,  $BC = \sqrt{5}$

$$\cos \angle ACB = \frac{AC^2 + BC^2 - AB^2}{2 \times AC \times BC}$$

$$= \frac{5 + 5 - 18}{2 \times \sqrt{5} \times \sqrt{5}}$$

$$= -\frac{8}{10} = -\frac{4}{5}$$

b  $1\frac{1}{2}$  cm<sup>2</sup>

8 a  $4$  b  $\frac{15\sqrt{3}}{4}$  (6.50) cm<sup>2</sup>

9 a  $1.50$  km b  $241^\circ$  c  $0.789$  km<sup>2</sup>

10  $359$  m<sup>2</sup>


11  $35.2$  m

12 a A stretch of scale factor 2 in the  $x$  direction.

b A translation of +3 in the  $y$  direction.




- c A reflection in the  $x$ -axis.  
 d A translation of  $-20$  in the  $x$  direction.



b There are no solutions.


- 14 a 300      b (30, 1)      c 60      d  $\frac{\sqrt{3}}{2}$

- 15 a  $p = 5$





b  $72^\circ$


- 16 a The four shaded regions are congruent.



- b  $\sin \alpha$  and  $\sin(180^\circ - \alpha)$  have the same  $y$  value, (call it  $k$ )  
 so  $\sin \alpha = \sin(180^\circ - \alpha)$   
 $\sin(180^\circ - \alpha)$  and  $\sin(360^\circ - \alpha)$  have the same  $y$  value, (which will be  $-k$ )  
 so  $\sin \alpha = \sin(180^\circ - \alpha)$   
 $= -\sin(180^\circ + \alpha)$   
 $= -\sin(360^\circ - \alpha)$



- b i From the graph of  $y = \cos \theta$ , which shows four congruent shaded regions, if the  $y$  value at  $\alpha$  is  $k$ , then  $y$  at  $180^\circ - \alpha$  is  $-k$ ,  $y$  at  $180^\circ - \alpha = -k$  and  $y$  at  $360^\circ - \alpha = +k$   
 so  $\cos \alpha = -\cos(180^\circ - \alpha)$   
 $= -\cos(180^\circ + \alpha)$   
 $= \cos(360^\circ - \alpha)$   
 ii From the graph of  $y = \tan \theta$ , if the  $y$  value at  $\alpha$  is  $k$ , then at  $180^\circ - \alpha$  it is  $-k$ , at  $180^\circ + \alpha$  it is  $+k$  and at  $360^\circ - \alpha$  it is  $-k$ ,  
 so  $\tan \alpha = -\tan(180^\circ - \alpha)$   
 $= +\tan(180^\circ + \alpha)$   
 $= -\tan(360^\circ - \alpha)$



- b 4  
 c The dunes may not all be the same height.

**Challenge**

Using the sine rule:


$$\sin(180^\circ - \angle ADB - \angle AEB) = \frac{5\left(\frac{1}{\sqrt{5}}\right)}{\sqrt{10}} = \frac{1}{\sqrt{2}}$$

$$180^\circ - \angle ADB - \angle AEB = 135^\circ \text{ (obtuse)}$$

$$\text{so } \angle ADB + \angle B = 45^\circ = \angle ACB$$


**CHAPTER 10**

**Prior knowledge check**




- b 4  
 c  $143.1^\circ, 396.9^\circ, 503.1^\circ$   
 2 a  $57.7^\circ$       b  $73.0^\circ$   
 3 a  $x = 11$       b  $x = \frac{9}{4}$       c  $x = -44.2^\circ$   
 4 a  $x = 1$  or  $x = 3$   
 b  $x = 1$  or  $x = -9$   
 c  $x = \frac{3 \pm \sqrt{65}}{4}$

**Exercise 10A**



- 2 a First      b Second      c Second  
 d Third      e Third
- 3 a -1      b 1      c 0      d -1      e -1  
 f 0      g 0      h 0      i 0      j 0
- 4 a  $-\sin 60^\circ$       b  $-\sin 80^\circ$       c  $\sin 20^\circ$   
 d  $-\sin 60^\circ$       e  $\sin 80^\circ$       f  $-\cos 70^\circ$   
 g  $-\cos 80^\circ$       h  $\cos 50^\circ$       i  $-\cos 20^\circ$   
 j  $-\cos 5^\circ$       k  $-\tan 80^\circ$       l  $-\tan 35^\circ$   
 m  $-\tan 30^\circ$       n  $\tan 5^\circ$       o  $\tan 60^\circ$
- 5 a  $-\sin \theta$       b  $-\sin \theta$       c  $-\sin \theta$   
 d  $\sin \theta$       e  $-\sin \theta$       f  $\sin \theta$   
 g  $-\sin \theta$       h  $-\sin \theta$       i  $\sin \theta$   
 j  $-\sin \theta$       k  $-\sin \theta$       l  $-\sin \theta$
- 6 a  $-\cos \theta$       b  $-\cos \theta$       c  $\cos \theta$   
 d  $-\cos \theta$       e  $\cos \theta$       f  $-\cos \theta$   
 g  $-\tan \theta$       h  $-\tan \theta$       i  $\tan \theta$   
 j  $\tan \theta$       k  $-\tan \theta$       l  $\tan \theta$

**Challenge**



$\sin \theta = \sin(180^\circ - \theta) = a$

$\cos \theta = \cos(-\theta) = b$

$\tan \theta = \frac{a}{b}; \tan(180^\circ - \theta) = \frac{a}{-b} = -\tan \theta$

For  $\tan \theta = \frac{x}{y}$

**Exercise 10B**

- 1 a  $\frac{\sqrt{2}}{2}$       b  $-\frac{\sqrt{3}}{2}$       c  $\frac{1}{2}$       d  $\frac{\sqrt{3}}{2}$   
 e  $\frac{\sqrt{3}}{2}$       f  $-\frac{1}{2}$       g  $\frac{1}{2}$       h  $-\frac{\sqrt{2}}{2}$   
 i  $-\frac{\sqrt{3}}{2}$       j  $-\frac{\sqrt{2}}{2}$       k -1      l -1  
 m  $\frac{\sqrt{3}}{3}$       n  $-\sqrt{3}$       o  $\sqrt{3}$

**Challenge**

- a i  $\sqrt{3}$       ii 2      iii  $\sqrt{2 + \sqrt{3}}$       iv  $\sqrt{2 + \sqrt{3}} - \sqrt{2}$   
 b  $15^\circ$   
 c i  $\frac{\sqrt{2 + \sqrt{3}} - \sqrt{2}}{2}$       ii  $\frac{\sqrt{2 + \sqrt{3}}}{2}$





**Exercise 10C**

- 1 a  $\sin^2 \frac{\theta}{2}$       b 5      c  $-\cos^2 A$   
 d  $\cos \theta$       e  $\tan x$       f  $\tan 3A$   
 g 4      h  $\sin^2 \theta$       i 1
- 2  $1\frac{1}{2}$
- 3  $\tan x - 3 \tan y$
- 4 a  $1 - \sin^2 \theta$       b  $\frac{\sin^2 \theta}{1 - \sin^2 \theta}$       c  $\sin \theta$   
 d  $\frac{1 - \sin^2 \theta}{\sin \theta}$       e  $1 - 2 \sin^2 \theta$
- 5 (One outline example of a proof is given)
- a LHS =  $\sin^2 \theta + \cos^2 \theta + 2 \sin \theta \cos \theta$   
 =  $1 + 2 \sin \theta \cos \theta$   
 = RHS
- b LHS =  $\frac{1 - \cos^2 \theta}{\cos \theta} = \frac{\sin^2 \theta}{\cos \theta} = \sin \theta \times \frac{\sin \theta}{\cos \theta}$   
 =  $\sin \theta \tan \theta = \text{RHS}$
- c LHS =  $\frac{\sin x}{\cos x} + \frac{\cos x}{\sin x} = \frac{\sin^2 x + \cos^2 x}{\sin x \cos x}$   
 =  $\frac{1}{\sin x \cos x} = \text{RHS}$
- d LHS =  $\cos^2 A - (1 - \cos^2 A) = 2 \cos^2 A - 1$   
 =  $2(1 - \sin^2 A) - 1 = 1 - 2 \sin^2 A = \text{RHS}$
- e LHS =  $(4 \sin^2 \theta - 4 \sin \theta \cos \theta + \cos^2 \theta) + (\sin^2 \theta + 4 \sin \theta \cos \theta + \cos^2 \theta)$   
 =  $5(\sin^2 \theta + \cos^2 \theta) = 5 = \text{RHS}$
- f LHS =  $2 - (\sin^2 \theta - 2 \sin \theta \cos \theta + \cos^2 \theta)$   
 =  $2(\sin^2 \theta + \cos^2 \theta) - (\sin^2 \theta - 2 \sin \theta \cos \theta + \cos^2 \theta)$   
 =  $\sin^2 \theta + 2 \sin \theta \cos \theta + \cos^2 \theta = (\sin \theta + \cos \theta)^2 = \text{RHS}$
- g LHS =  $\sin^2 x (1 - \sin^2 y) - (1 - \sin^2 x) \sin^2 y$   
 =  $\sin^2 x - \sin^2 y = \text{RHS}$
- 6 a  $\sin \theta = \frac{5}{13}, \cos \theta = \frac{12}{13}$   
 b  $\sin \theta = \frac{4}{5}, \tan \theta = -\frac{4}{3}$   
 c  $\cos \theta = \frac{24}{25}, \tan \theta = -\frac{7}{24}$
- 7 a  $-\frac{\sqrt{5}}{3}$       b  $\frac{2\sqrt{5}}{5}$
- 8 a  $-\frac{\sqrt{3}}{2}$       b  $\frac{1}{2}$
- 9 a  $\frac{\sqrt{7}}{4}$       b  $-\frac{\sqrt{7}}{3}$
- 10 a  $x^2 + y^2 = 1$   
 b  $4x^2 + y^2 = 4$       (or  $x^2 + \frac{y^2}{4} = 1$ )  
 c  $x^2 + y = 1$   
 d  $x^2 = y^2 (1 - x^2)$       (or  $x^2 + \frac{x^2}{y^2} = 1$ )  
 e  $x^2 + y^2 = 2$       (or  $\frac{(x+y)^2}{4} + \frac{(x-y)^2}{4} = 1$ )
- 11 a Using cosine rule:  $\cos B = \frac{8^2 + 12^2 - 10^2}{2 \times 8 \times 12} = \frac{9}{16}$   
 b  $\frac{\sqrt{175}}{16}$
- 12 a Using sine rule:  $\sin Q = \frac{\sin 30}{6} \times 8 = \frac{2}{3}$   
 b  $-\frac{\sqrt{5}}{3}$

**Exercise 10D**

- 1 a  $-63.4^\circ$       b  $116.6^\circ, 296.6^\circ$   
 2 a  $66.4^\circ$       b  $66.4^\circ, 113.6^\circ, 246.4^\circ, 293.6^\circ$   
 3 a  $270^\circ$       b  $60^\circ, 240^\circ$   
 c  $60^\circ, 300^\circ$       d  $15^\circ, 165^\circ$   
 e  $140^\circ, 220^\circ$       f  $135^\circ, 315^\circ$   
 g  $90^\circ, 270^\circ$       h  $230^\circ, 310^\circ$
- 4 a  $45.6^\circ, 134.4^\circ$       b  $135^\circ, 225^\circ$   
 c  $132^\circ, 228^\circ$       d  $229^\circ, 311^\circ$   
 e  $8.13^\circ, 188^\circ$       f  $61.9^\circ, 242^\circ$   
 g  $105^\circ, 285^\circ$       h  $41.8^\circ, 318^\circ$
- 5 a  $30^\circ, 210^\circ$       b  $135^\circ, 315^\circ$   
 c  $53.1^\circ, 233^\circ$       d  $56.3^\circ, 236^\circ$   
 e  $54.7^\circ, 235^\circ$       f  $148^\circ, 328^\circ$
- 6 a  $-120^\circ, -60^\circ, 240^\circ, 300^\circ$       b  $-171^\circ, -8.63^\circ$   
 c  $-144^\circ, 144^\circ$       d  $-327^\circ, -32.9^\circ$   
 e  $150^\circ, 330^\circ, 510^\circ, 690^\circ$       f  $251^\circ, 431^\circ$
- 7 a  $\tan x$  should be  $\frac{2}{3}$   
 b Squaring both sides creates extra solutions  
 c  $-146.3^\circ, 33.7^\circ$
- 8 a
- b 2      c  $26.6^\circ, 206.6^\circ$
- 9  $71.6^\circ, 108.4^\circ, 251.6^\circ, 288.4^\circ$
- 10 a  $4 \sin^2 x - 3(1 - \sin^2 x) = 2$ .  
 Rearrange to get  $7 \sin^2 x = 5$   
 b  $57.7^\circ, 122.3^\circ, 237.7^\circ, 302.3^\circ$
- 11 a  $2 \sin^2 x + 5(1 - \sin^2 x) = 1$ .  
 Rearrange to get  $3 \sin^2 x = 4$   
 b  $\sin x > 1$

**Exercise 10E**

- 1 a  $0^\circ, 45^\circ, 90^\circ, 135^\circ, 180^\circ, 225^\circ, 270^\circ, 315^\circ, 360^\circ$   
 b  $60^\circ, 180^\circ, 300^\circ$   
 c  $22\frac{1}{2}^\circ, 112\frac{1}{2}^\circ, 202\frac{1}{2}^\circ, 292\frac{1}{2}^\circ$   
 d  $30^\circ, 150^\circ, 210^\circ, 330^\circ$   
 e  $300^\circ$   
 f  $225^\circ, 315^\circ$
- 2 a  $90^\circ, 270^\circ$       b  $50^\circ, 170^\circ$       c  $165^\circ, 345^\circ$   
 d  $250^\circ, 310^\circ$       e  $16.9^\circ, 123^\circ$
- 3 a  $11.2^\circ, 71.2^\circ, 131.2^\circ$       b  $6.3^\circ, 186.3^\circ, 366.3^\circ$   
 c  $37.0^\circ, 127.0^\circ$       d  $-150^\circ, 30^\circ$
- 4 a  $10^\circ, 130^\circ$       b  $71.6^\circ, 108.4^\circ$
- 5 a

- b  $(0^\circ, \frac{\sqrt{3}}{2}), (120^\circ, 0), (300^\circ, 0)$   
 c  $86.6^\circ, 333.4^\circ$   
 6 a  $0.75$   
 b  $18.4^\circ, 108.4^\circ, 198.4^\circ, 288.4^\circ$   
 7 a  $2.5$   
 b No: increasing  $k$  will bring another 'branch' of the tan graph into place.

**Challenge**  
 $25^\circ, 65^\circ, 145^\circ$

**Exercise 10F**


- 1 a  $60^\circ, 120^\circ, 240^\circ, 300^\circ$   
 b  $45^\circ, 135^\circ, 225^\circ, 315^\circ$   
 c  $0^\circ, 180^\circ, 199^\circ, 341^\circ, 360^\circ$   
 d  $77.0^\circ, 113^\circ, 257^\circ, 293^\circ$   
 e  $60^\circ, 300^\circ$   
 f  $204^\circ, 336^\circ$   
 g  $30^\circ, 60^\circ, 120^\circ, 150^\circ, 210^\circ, 240^\circ, 300^\circ, 330^\circ$   
 2 a  $\pm 45^\circ, \pm 135^\circ$  b  $-180^\circ, -117^\circ, 0^\circ, 63.4^\circ, 180^\circ$   
 c  $\pm 114^\circ$  d  $0^\circ, \pm 75.5^\circ, \pm 180^\circ$   
 3 a  $72.0^\circ, 144^\circ$  b  $0^\circ, 60^\circ$   
 c No solutions in range  
 4 a  $\pm 41.8^\circ, \pm 138^\circ$  b  $38.2^\circ, 142^\circ$   
 5  $60^\circ, 75.5^\circ, 284.5^\circ, 300^\circ$   
 6  $48.2^\circ, 131.8^\circ, 228.2^\circ, 311.8^\circ$   
 7  $2 \cos^2 x + \cos x - 6 = (2 \cos x - 3)(\cos x + 2)$   
 There are no solutions to  $\cos x = -2$  or to  $\cos x = \frac{3}{2}$   
 8 a  $1 - \sin^2 x = 2 - \sin x$   
 Rearrange to get  $\sin^2 x - \sin x + 1 = 0$   
 b The equation has no real roots as  $b^2 - 4ac < 0$   
 9 a  $p = 1, q = 5$   
 b  $72.8^\circ, 129.0^\circ, 252.8^\circ, 309.0^\circ, 432.8^\circ, 489.0^\circ$

**Challenge**  
 1  $-180^\circ, -60^\circ, 60^\circ, 180^\circ$   
 2  $0^\circ, 90^\circ, 180^\circ, 270^\circ, 360^\circ$

**Mixed Exercise**

- 1 a  $-\cos 57^\circ$  b  $-\sin 48^\circ$  c  $+\tan 10^\circ$   
 2 a  $0$  b  $-\frac{\sqrt{2}}{2}$  c  $-1$   
 d  $\sqrt{3}$  e  $-1$   
 3 Using  $\sin^2 A = 1 - \cos^2 A$ ,  $\sin^2 A = 1 - \left(-\sqrt{\frac{7}{11}}\right)^2 = \frac{4}{11}$ .  
 Since angle  $A$  is obtuse, it is in the second quadrant and  $\sin$  is positive, so  $\sin A = \frac{2}{\sqrt{11}}$ .  
 Then  $\tan A = \frac{\sin A}{\cos A} = \frac{2}{\sqrt{11}} \times \left(-\sqrt{\frac{11}{7}}\right) = -\frac{2}{\sqrt{7}} = -\frac{2}{7}\sqrt{7}$ .  
 4 a  $-\frac{\sqrt{21}}{5}$  b  $-\frac{2}{5}$   
 5 a  $\cos^2 \theta - \sin^2 \theta$  b  $\sin^4 3\theta$  c  $1$   
 6 a  $1$  b  $\tan y = \frac{4 + \tan x}{2 \tan x - 3}$   
 7 a LHS  $= (1 + 2 \sin \theta + \sin^2 \theta) + \cos^2 \theta$   
 $= 1 + 2 \sin \theta + 1$   
 $= 2 + 2 \sin \theta$   
 $= 2(1 + \sin \theta) = \text{RHS}$

- b LHS  $= \cos^4 \theta + \sin^2 \theta$   
 $= (1 - \sin^2 \theta)^2 + \sin^2 \theta$   
 $= 1 - 2 \sin^2 \theta + \sin^4 \theta + \sin^2 \theta$   
 $= (1 - \sin^2 \theta) + \sin^4 \theta$   
 $= \cos^2 \theta + \sin^4 \theta = \text{RHS}$   
 8 a No solutions:  $-1 \leq \sin \theta \leq 1$   
 b 2 solutions:  $\tan \theta = -1$  has two solutions in the interval.  
 c No solutions:  $2 \sin \theta + 3 \cos \theta > -5$   
 so  $2 \sin \theta + 3 \cos \theta + 6$  can never be equal to 0.  
 d No solutions:  $\tan^2 \theta = -1$  has no real solutions.  
 9 a  $(4x - y)(y + 1)$  b  $14.0^\circ, 180^\circ, 194^\circ$   
 10 a  $3 \cos 3\theta$  b  $16.1, 104, 136, 224, 256, 344$   
 11 a  $2 \sin 2\theta = \cos 2\theta \Rightarrow \frac{2 \sin 2\theta}{\cos 2\theta} = 1$   
 $\Rightarrow 2 \tan 2\theta = 1 \Rightarrow \tan 2\theta = 0.5$   
 b  $13.3^\circ, 103.3^\circ, 193.3^\circ, 283.3^\circ$   
 12 a  $225^\circ, 345^\circ$   
 b  $22.2^\circ, 67.8^\circ, 202.2^\circ, 247.8^\circ$   
 13  $30^\circ, 150^\circ, 210^\circ$   
 14  $0^\circ, 131.8^\circ, 228.2^\circ$   
 15 a Found additional solutions after dividing by three rather than before. Not applied the full interval for solutions.  
 b  $-350^\circ, -310^\circ, -230^\circ, -190^\circ, -110^\circ, -70^\circ, 10^\circ, 50^\circ, 130^\circ, 170^\circ, 250^\circ, 290^\circ$




- b  $2$  c  $33.7^\circ, 213.7^\circ$   
 17 a  $\frac{9}{11}$  b  $\frac{\sqrt{40}}{11}$   
 18 a Using sine rule:  $\sin Q = \sin 45 \times \frac{6}{5} = \frac{\sqrt{2}}{2} \times \frac{6}{5} = \frac{3\sqrt{2}}{5}$   
 b  $\frac{\sqrt{7}}{5}$   
 19 a  $3 \sin^2 x - (1 - \sin^2 x) = 2$ .  
 Rearrange to give  $4 \sin^2 x = 3$ .  
 b  $-120^\circ, -60^\circ, 60^\circ, 120^\circ$   
 20  $-318.2^\circ, -221.8^\circ, 41.8^\circ, 138.2^\circ$


**Challenge**  
 $45^\circ, 54.7^\circ, 125.3^\circ, 135^\circ, 225^\circ, 234.7^\circ, 305.3^\circ, 315^\circ$

**Review exercise 2**

- 1  $x + 3y - 22 = 0$   
 2  $x - 3y - 21 = 0$   
 3  $4, -2.5$   
 4 a  $0.45$   
 b  $l = 0.45h$   
 c The model may not be valid for young people who are still growing.  
 5 a  $y = -\frac{1}{3}x + 4$  b C is  $(3, 3)$  c  $15$



- 6  $3\sqrt{5}$   
 7  $(-6, 0)$   
 8  $(x + 3)^2 + (y - 8)^2 = 10$   
 9 a  $(x - 3)^2 + (y + 1)^2 = 20$  ( $a = 3, b = -1, r = \sqrt{20}$ )  
 b Centre  $(3, -1)$ , radius  $\sqrt{20}$   
 10 a  $(3, 5)$  and  $(4, 2)$   
 b  $\sqrt{10}$   
 11  $0 < r < \sqrt{\frac{2}{5}}$   
 12 a  $(x - 1)^2 + (y - 5)^2 = 58$   
 b  $7y - 3x + 26 = 0$   
 13 a  $AB = \sqrt{32}; BC = \sqrt{8}; AC = \sqrt{40}; AC^2 = AB^2 + BC^2$   
 b  $AC$  is a diameter of the circle.  
 c  $(x - 5)^2 + (y - 2)^2 = 10$   
 14  $a = 3, b = -2, c = -8$   
 15 a  $2(\frac{1}{2})^3 - 7(\frac{1}{2})^2 - 17(\frac{1}{2}) + 10 = 0$   
 b  $(2x - 1)(x - 5)(x + 2)$   
 c
- 
- 16 a 24  
 b  $(x - 3)(3x - 2)(x + 4)$   
 17 a  $g(3) = 3^3 - 13(3) + 12 = 0$   
 b  $(x - 3)(x + 4)(x - 1)$   
 18 a  $a = 0, b = 0$   
 b  $a > 0, b > 0$   
 19 a  $5^2 = 24 + 1; 7^2 = 2(24) + 1; 11^2 = 5(24) + 1;$   
 $13^2 = 7(24) + 1; 17^2 = 12(24) + 1; 19^2 = 15(24) + 1$   
 b  $3(24) + 1 = 73$  which is not a square of a prime number  
 20 a  $(x - 5)^2 + (y - 4)^2 = 3^2$   
 b  $\sqrt{41}$   
 c Sum of radii  $= 3 + 3 < \sqrt{41}$  so circles do not touch  
 21 a  $1 - 20x + 180x^2 - 960x^3 + \dots$   
 b 0.817  
 22  $a = 2, b = 19, c = 70$   
 23 4  
 24  $\sqrt{10}$  cm  
 25 a  $\cos 60^\circ = \frac{1}{2} = \frac{(5^2 + (2x - 3)^2 - (x + 1)^2) \div 2(5)(2x - 3)}{5(2x - 3)} = \frac{(25 + 4x^2 - 12x + 9 - x^2 - 2x - 1) \div 2(5)(2x - 3)}{5(2x - 3)}$   
 $0 = 3x^2 - 24x + 48$   
 $x^2 - 8x + 16 = 0$   
 b 4  
 c  $10.8 \text{ cm}^2$   
 26 a 11.93 km  
 b  $100.9^\circ$   
 27 a  $AB = BC = 10 \text{ cm}, AC = 6\sqrt{10} \text{ cm}$   
 b  $143.1^\circ$   
 28  $19.4 \text{ km}^2$   
 29 a  $(x - 5)^2 + (y - 2)^2 = 25$   
 b 6  
 c  $XY = \sqrt{90}; YZ = \sqrt{20}; XZ = \sqrt{98}$   
 $\cos XYZ = (20 + 90 - 98) \div (2 \times \sqrt{20} \times \sqrt{90})$   
 $\cos XYZ = 12 \div 60\sqrt{2} = \sqrt{2} \div 10$



- b 2  
 31 a  $(-225, 0), (-45, 0), (135, 0)$  and  $(315, 0)$   
 b  $(0, \frac{\sqrt{2}}{2})$   
 32 Area of triangle  $= \frac{1}{2} \times s \times s \times \sin 60^\circ = \frac{\sqrt{3}}{4} s^2$   
 Area of square  $= s^2$   
 Total surface area  $= 4 \times (\frac{\sqrt{3}}{4} s^2) + s^2 = (\sqrt{3} + 1)s^2 \text{ cm}^2$   
 33 a 1  
 b  $45^\circ, 225^\circ$   
 34  $30^\circ, 150^\circ, 210^\circ, 330^\circ$   
 35  $90^\circ, 150^\circ$   
 36 a  $2(1 - \sin^2 x) = 4 - 5 \sin x$   
 $2 - 2 \sin^2 x = 4 - 5 \sin x$   
 $2 \sin^2 x - 5 \sin x + 2 = 0$   
 b  $x = 30^\circ, 150^\circ$   
 37  $72.3^\circ, 147.5^\circ, 252.3^\circ, 327.5^\circ$   
 38  $0^\circ, 78.5^\circ, 281.5^\circ, 360^\circ$   
 39  $\cos^2 x (\tan^2 x + 1)$   
 $= \cos^2 x (\frac{\sin^2 x}{\cos^2 x} + 1)$   
 $= \cos^2 x + \sin^2 x = 1$

Challenge

- 1 a 160  
 b  $(-\frac{28}{3}, 0)$   
 2 The second circle has the same centre but a larger radius  
 3  $\binom{n}{k} + \binom{n}{k+1} = \frac{n!}{k!(n-k)!} + \frac{n!}{(k+1)!(n-k-1)!}$   
 $= \frac{n!(k+1)}{(k+1)!(n-k)!} + \frac{n!(n-k)}{(k+1)!(n-k)!}$   
 $= \frac{n!((k+1) + (n-k))}{(k+1)!(n-k)!}$   
 $= \frac{n!(n+1)}{(k+1)!(n-k)!}$   
 $= \frac{(n+1)!}{(k+1)!(n-k)!}$   
 $= \binom{n+1}{k+1}$   
 4  $0^\circ, 30^\circ, 150^\circ, 180^\circ, 270^\circ, 360^\circ$


CHAPTER 11

Prior knowledge check

- 1 a  $\begin{pmatrix} 4 \\ 2 \end{pmatrix}$       b  $\begin{pmatrix} 5 \\ 2 \end{pmatrix}$       c  $\begin{pmatrix} -1 \\ -3 \end{pmatrix}$   
 2 a  $\frac{7}{9}$       b  $\frac{2}{9}$       c  $\frac{7}{2}$   
 3 a  $123.2^\circ$       b 13.6      c 5.3      d  $21.4^\circ$



**Exercise 11A**



- 2 a  $2b$       b  $d$       c  $b$   
 d  $2b$       e  $d + b$       f  $d + b$   
 g  $-2d$       h  $-b$       i  $2d + b$   
 j  $-b + 2d$       k  $-b + d$       l  $-d - b$
- 3 a  $2m$       b  $2p$       c  $m$   
 d  $m$       e  $p + m$       f  $p + m$   
 g  $p + 2m$       h  $p - m$       i  $-m - p$   
 j  $-2m + p$       k  $-2p + m$       l  $-m - 2p$
- 4 a  $d - a$       b  $a + b + c$   
 c  $a + b - d$       d  $a + b + c - d$
- 5 a  $2a + 2b$       b  $a + b$       c  $b - a$   
 6 a  $b$       b  $b - 3a$       c  $a - b$   
 d  $2a - b$
- 7 a  $\vec{OB} = a + b$       b  $\vec{OP} = \frac{5}{8}(a + b)$       c  $\vec{AP} = \frac{5}{8}b - \frac{3}{8}a$
- 8 a Yes ( $\lambda = 2$ )      b Yes ( $\lambda = 4$ )      c No  
 d Yes ( $\lambda = -1$ )      e Yes ( $\lambda = -3$ )      f No
- 9 a i  $b - a$       ii  $\frac{1}{2}a$       iii  $\frac{1}{2}b$       iv  $\frac{1}{2}b - \frac{1}{2}a$   
 b  $\vec{BC} = b - a$ ,  $\vec{PQ} = \frac{1}{2}(b - a)$  so  $PQ$  is parallel to  $BC$ .
- 10 a i  $2b$       ii  $a - b$   
 b  $\vec{AB} = 2b$ ,  $\vec{OC} = 3b$  so  $AB$  is parallel to  $OC$ .
- 11 1.2


**Exercise 11B**

- 1  $v_1: 8i, \begin{pmatrix} 8 \\ 0 \end{pmatrix}$        $v_2: 9i + 3j, \begin{pmatrix} 9 \\ 3 \end{pmatrix}$        $v_3: -4i + 2j, \begin{pmatrix} -4 \\ 2 \end{pmatrix}$   
 $v_4: 3i + 5j, \begin{pmatrix} 3 \\ 5 \end{pmatrix}$        $v_5: -3i - 2j, \begin{pmatrix} -3 \\ -2 \end{pmatrix}$        $v_6: -5j, \begin{pmatrix} 0 \\ -5 \end{pmatrix}$

- 2 a  $8i + 12j$       b  $i + 1.5j$       c  $-4i + j$   
 d  $10i + j$       e  $-2i + 11j$       f  $-2i - 10j$   
 g  $14i - 7j$       h  $-8i + 9j$
- 3 a  $\begin{pmatrix} 45 \\ 35 \end{pmatrix}$       b  $\begin{pmatrix} 4 \\ 0.5 \end{pmatrix}$       c  $\begin{pmatrix} 12 \\ 3 \end{pmatrix}$   
 d  $\begin{pmatrix} -1 \\ 16 \end{pmatrix}$       e  $\begin{pmatrix} -21 \\ -29 \end{pmatrix}$       f  $\begin{pmatrix} 10 \\ 2 \end{pmatrix}$
- 4 a  $\lambda = 5$       b  $\mu = -\frac{3}{2}$   
 5 a  $\lambda = \frac{1}{3}$       b  $\mu = -1$   
 c  $s = -1$       d  $t = -\frac{1}{17}$
- 6  $i - j$
- 7 a  $\vec{AC} = 5i - 4j = \begin{pmatrix} 5 \\ -4 \end{pmatrix}$       b  $\vec{OP} = 5i + \frac{8}{5}j = \begin{pmatrix} 5 \\ \frac{8}{5} \end{pmatrix}$   
 c  $\vec{AP} = 3i - \frac{12}{5}j = \begin{pmatrix} 3 \\ -\frac{12}{5} \end{pmatrix}$
- 8  $j = 4, k = 11$   
 9  $p = 3, q = 2$   
 10 a  $p = 5$       b  $8i - 12j$

**Exercise 11C**


- 1 a 5      b 10      c 13  
 d 4.47 (3 s.f.)      e 5.83 (3 s.f.)      f 8.06 (3 s.f.)  
 g 5.83 (3 s.f.)      h 4.12 (3 s.f.)
- 2 a  $\sqrt{26}$       b  $5\sqrt{2}$       c  $\sqrt{101}$
- 3 a  $\frac{1}{5} \begin{pmatrix} 4 \\ 3 \end{pmatrix}$       b  $\frac{1}{13} \begin{pmatrix} 5 \\ -12 \end{pmatrix}$   
 c  $\frac{1}{25} \begin{pmatrix} -7 \\ 24 \end{pmatrix}$       d  $\frac{1}{\sqrt{10}} \begin{pmatrix} 1 \\ -3 \end{pmatrix}$
- 4 a  $53.1^\circ$  above      b  $53.1^\circ$  below  
 c  $67.4^\circ$  above      d  $63.4^\circ$  above
- 5 a  $149^\circ$  to the right      b  $29.7^\circ$  to the right  
 c  $31.0^\circ$  to the left      d  $104^\circ$  to the left
- 6 a  $\frac{15\sqrt{2}}{2}i + \frac{15\sqrt{2}}{2}j, \begin{pmatrix} 15\sqrt{2} \\ 2 \\ 15\sqrt{2} \\ 2 \end{pmatrix}$       b  $7.52i + 2.74j, \begin{pmatrix} 7.52 \\ 2.74 \end{pmatrix}$   
 c  $18.1i - 8.45j, \begin{pmatrix} 18.1 \\ -8.45 \end{pmatrix}$       d  $\frac{5\sqrt{3}}{2}i - 2.5j, \begin{pmatrix} 5\sqrt{3} \\ 2 \\ -2.5 \end{pmatrix}$
- 7 a  $|3i + 4j| = 5, 53.1^\circ$  above



b  $|2i - j| = \sqrt{5}, 26.6^\circ$  below



c  $|-5i + 2j| = \sqrt{29}, 158.2^\circ$  above



- 8  $k = \pm 6$   
 9  $p = \pm 8, q = 6$






- 10 a  $36.9^\circ$       b  $33.7^\circ$       c  $70.6^\circ$   
 11 a  $67.2^\circ$       b  $19.0$

**Challenge**

Possible solution:



Area of parallelogram = area of large rectangle – 2(area of small rectangle) – 2(area triangle 1) – 2(area triangle 2)

$$\begin{aligned} \text{Area of parallelogram} &= (p+r)(q+s) - 2qr - 2\left(\frac{1}{2}pq\right) \\ &\quad - 2\left(\frac{1}{2}rs\right) = ps - qr \end{aligned}$$

**Exercise 11D**

- 1 a i  $\vec{OA} = 3\mathbf{i} - \mathbf{j}$ ,  $\vec{OB} = 4\mathbf{i} + 5\mathbf{j}$ ,  $\vec{OC} = -2\mathbf{i} + 6\mathbf{j}$   
 ii  $\mathbf{i} + 6\mathbf{j}$       iii  $-5\mathbf{i} + 7\mathbf{j}$   
 b i  $\sqrt{40} = 2\sqrt{10}$       ii  $\sqrt{37}$       iii  $\sqrt{74}$
- 2 a  $-\mathbf{i} + 5\mathbf{j}$  or  $\begin{pmatrix} -1 \\ 5 \end{pmatrix}$   
 b i 5      ii  $\sqrt{13}$       iii  $\sqrt{26}$
- 3 a  $-\mathbf{i} - 9\mathbf{j}$  or  $\begin{pmatrix} -1 \\ -9 \end{pmatrix}$   
 b i  $\sqrt{82}$       ii 5      iii  $\sqrt{61}$
- 4 a  $-2\mathbf{a} + 2\mathbf{b}$       b  $-3\mathbf{a} + 2\mathbf{b}$       c  $-2\mathbf{a} + \mathbf{b}$
- 5  $\begin{pmatrix} 7 \\ 9 \end{pmatrix}$  or  $\begin{pmatrix} 9 \\ 3 \end{pmatrix}$
- 6 a  $2\mathbf{i} + 8\mathbf{j}$       b  $2\sqrt{17}$
- 7  $\frac{3\sqrt{5}}{5}$

**Challenge**

$$\vec{OB} = 2\mathbf{i} + 3\mathbf{j} \text{ or } \vec{OB} = \frac{46}{13}\mathbf{i} + \frac{9}{13}\mathbf{j}$$

**Exercise 11E**

- 1  $\vec{XY} = \mathbf{b} - \mathbf{a}$  and  $\vec{YZ} = \mathbf{c} - \mathbf{b}$ , so  $\mathbf{b} - \mathbf{a} = \mathbf{c} - \mathbf{b}$ .  
 Hence  $\mathbf{a} + \mathbf{c} = 2\mathbf{b}$ .
- 2 a i  $2\mathbf{r}$       ii  $\mathbf{r}$   
 b Sides of triangle  $OAB$  are twice the length of sides of triangle  $PAQ$  and angle  $A$  is common to both  $SAS$ .
- 3 a  $\frac{2}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}$   
 b  $\vec{AN} = \frac{1}{3}(\mathbf{b} - \mathbf{a})$ ,  $\vec{AB} = \mathbf{b} - \mathbf{a}$ ,  $\vec{NB} = \frac{2}{3}(\mathbf{b} - \mathbf{a})$   
 so  $AN:NB = 1:2$ .
- 4 a  $\frac{3}{5}\mathbf{a} + \frac{2}{5}\mathbf{c}$   
 b  $\vec{AP} = -\mathbf{a} + \frac{3}{5}\mathbf{a} + \frac{2}{5}\mathbf{c} = \frac{2}{5}(\mathbf{c} - \mathbf{a})$ ,  
 $\vec{PC} = \mathbf{c} - (\frac{3}{5}\mathbf{a} + \frac{2}{5}\mathbf{c}) = \frac{3}{5}(\mathbf{c} - \mathbf{a})$  so  $AP:PC = 2:3$
- 5 a  $\sqrt{26}$       b  $2\sqrt{2}$       c  $3\sqrt{2}$   
 d  $\angle BAC = 56^\circ$ ,  $\angle ABC = 34^\circ$ ,  $\angle ACB = 90^\circ$
- 6 a  $\vec{OR} = \mathbf{a} + \frac{1}{3}(\mathbf{b} - \mathbf{a}) = \frac{2}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}$ ,  
 $\vec{OS} = 3\vec{OR} = 3(\frac{2}{3}\mathbf{a} + \frac{1}{3}\mathbf{b}) = 2\mathbf{a} + \mathbf{b}$

- b  $\vec{TP} = \mathbf{a} + \mathbf{b}$ ,  $\vec{PS} = \frac{1}{3}(\mathbf{b} - \mathbf{a}) + \frac{2}{3}(2\mathbf{a} + \mathbf{b}) = \mathbf{a} + \mathbf{b}$   
 $\vec{TP}$  is parallel (and equal) to  $\vec{PS}$  and they have a point,  $P$ , in common so  $T, P$  and  $S$  lie on a straight line.

**Challenge:**

- a  $\vec{PR} = \mathbf{b} - \mathbf{a}$ ,  $\vec{PX} = j(\mathbf{b} - \mathbf{a}) = -j\mathbf{a} + j\mathbf{b}$   
 b  $\vec{ON} = \mathbf{a} + \frac{1}{2}\mathbf{b}$ ,  $\vec{PX} = -\mathbf{a} + k(\mathbf{a} + \frac{1}{2}\mathbf{b}) = (k-1)\mathbf{a} + \frac{1}{2}k\mathbf{b}$   
 c Coefficients of  $\mathbf{a}$  and  $\mathbf{b}$  must be the same in both expressions for  $\vec{PX}$   
 Coefficients of  $\mathbf{a}$ :  $k-1 = -j$ ; Coefficients of  $\mathbf{b}$ :  $j = \frac{1}{2}k$   
 d Solving simultaneously gives  $j = \frac{1}{3}$  and  $k = \frac{2}{3}$   
 e  $\vec{PX} = \frac{1}{3}\vec{PR}$ .  
 By symmetry,  $\vec{PX} = \vec{YR} = \vec{XY}$ , so  $ON$  and  $OM$  divide  $PR$  into 3 equal parts.

**Exercise 11F**

- 1 a  $5 \text{ m s}^{-1}$       b  $25 \text{ km h}^{-1}$   
 c  $5.39 \text{ m s}^{-1}$       d  $8.06 \text{ cm s}^{-1}$
- 2 a 50 km      b 51.0 m  
 c 4.74 km      d 967 cm
- 3 a  $5 \text{ m s}^{-1}$ , 75 m      b  $5.39 \text{ m s}^{-1}$ , 16.2 m  
 c  $5.39 \text{ km h}^{-1}$ , 16.2 km      d  $13 \text{ km h}^{-1}$ , 6.5 km
- 4  $(2.8\mathbf{i} - 1.6\mathbf{j}) \text{ m s}^{-2}$
- 5 a  $54.5^\circ$       b  $0.3\sqrt{74}$  Newtons
- 6 a  $26.6^\circ$  below  $\mathbf{i}$   
 b  $\mathbf{R} = (3+p)\mathbf{i} + (q-4)\mathbf{j}$ ,  $3+p = 2\lambda$  and  $q-4 = -\lambda \Rightarrow \lambda = 4-q$   
 $3+p = 2(4-q) \Rightarrow 3+p = 8-2q$  so  $p+2q = 5$
- c  $|\mathbf{R}| = 2\sqrt{5}$  newtons
- 7 a  $10\mathbf{i} - 100\mathbf{j}$       b  $109.4^\circ$       c  $1700 \text{ m}^2$
- 8 a  $\sqrt{41}$       b  $303.7^\circ$   
 c  $\vec{AB} = 4\mathbf{i} - 5\mathbf{j}$ ,  $\mathbf{v} = 2(4\mathbf{i} - 5\mathbf{j})$  so the boat is travelling directly towards the buoy.  
 d  $2\sqrt{41}$       e 30 minutes

**Mixed exercise**

- 1 a  $2\sqrt{10}$  newtons      b  $18^\circ$   
 2 a  $108^\circ$       b  $9.48 \text{ km h}^{-1}$   
 3 a  $9.85 \text{ m s}^{-1}$       b 59.1 m  
 c The model ignores friction and air resistance.  
 The model will become less accurate as  $t$  increases.
- 4 a  $\mathbf{b} - \frac{3}{5}\mathbf{a}$       b  $\mathbf{b} - 4\mathbf{a}$       c  $\frac{2}{5}\mathbf{a} - \mathbf{b}$       d  $3\mathbf{a} - \mathbf{b}$
- 5 1.25
- 6 a  $\begin{pmatrix} 12 \\ -1 \end{pmatrix}$       b  $\begin{pmatrix} -18 \\ 5 \end{pmatrix}$       c  $\begin{pmatrix} 49 \\ 13 \end{pmatrix}$
- 7 a  $3\mathbf{i} - 2\mathbf{j}$       b  $32.5^\circ$       c 10.5
- 8 a  $p = -1.5$       b  $\mathbf{i} - 1.5\mathbf{j}$
- 9 a i  $\frac{1}{17}(8\mathbf{i} + 15\mathbf{j})$       ii  $61.9^\circ$  above  
 b i  $\frac{1}{25}(24\mathbf{i} - 7\mathbf{j})$       ii  $16.3^\circ$  below  
 c i  $\frac{1}{41}(-9\mathbf{i} + 40\mathbf{j})$       ii  $102.7^\circ$  above  
 d i  $\frac{1}{\sqrt{13}}(3\mathbf{i} - 2\mathbf{j})$       ii  $33.7^\circ$  below
- 10  $p = 8.6$ ,  $q = 12.3$
- 11  $\pm 6$
- 12 a  $\frac{3}{5}\mathbf{a} + \frac{2}{5}\mathbf{b}$       b  $\frac{2}{5}\mathbf{b}$   
 c  $\vec{AB} = \mathbf{b} - \mathbf{a}$ ,  $\vec{AN} = \frac{2}{5}(\mathbf{b} - \mathbf{a})$  so  $AN:NB = 2:3$

- 13 a  $18.4^\circ$  below  
 b  $\mathbf{R} = (4 + p)\mathbf{i} + (5 - q)\mathbf{j}$ ,  $4 + p = 3\lambda$  and  $5 - q = -\lambda$   
 $4 + p = 3(q - 5)$  so  $p + 3q = 11$   
 c  $2\sqrt{10}$  newtons  
 14  $\frac{\sqrt{193}}{2}$

**Challenge**

$\overrightarrow{OB} = \frac{3}{2}\mathbf{i} + \frac{5}{2}\mathbf{j}$  or  $\frac{99}{34}\mathbf{i} + \frac{5}{34}\mathbf{j}$

**CHAPTER 12**

**Prior knowledge check**

- 1 a 5      b  $-\frac{2}{3}$       c  $\frac{1}{3}$   
 2 a  $x^{10}$       b  $x^{\frac{2}{3}}$       c  $x^{-1}$       d  $x^{\frac{1}{4}}$   
 3 a  $y = \frac{1}{2}x - 2$       b  $y = -\frac{1}{2}x + 8\frac{1}{2}$       c  $y = -\frac{1}{4}x + 7\frac{1}{2}$   
 4  $y = -\frac{1}{2}x$  or  $y = -\frac{1}{3}x + 1\frac{2}{3}$

**Exercise 12A**

1 a	x-coordinate	-1	0	1	2	3
	Estimate for gradient of curve	-4	-2	0	2	4

- b Gradient =  $2p - 2$       c 1  
 2 a  $\sqrt{1 - 0.6^2} = \sqrt{0.64} = 0.8$   
 b Gradient =  $-0.75$   
 c i  $-1.21$  (3 s.f.)      ii  $-1$       iii  $-0.859$  (3 s.f.)  
 d As other point moves closer to A, gradient tends to  $-0.75$ .  
 3 a i 7      ii 6.5      iii 6.1  
 iv 6.01      v  $h + 6$   
 b 6  
 4 a i 9      ii 8.5      iii 8.1  
 iv 8.01      v  $8 + h$   
 b 8

**Exercise 12B**

- 1 a  $f'(2) = \lim_{h \rightarrow 0} \frac{f(2+h) - f(2)}{h} = \lim_{h \rightarrow 0} \frac{(2+h)^2 - 2^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{4h + h^2}{h} = \lim_{h \rightarrow 0} (4 + h) = 4$   
 b  $f'(-3) = \lim_{h \rightarrow 0} \frac{f(-3+h) - f(-3)}{h} = \lim_{h \rightarrow 0} \frac{(-3+h)^2 - 3^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{-6h + h^2}{h} = \lim_{h \rightarrow 0} (-6 + h) = -6$   
 c  $f'(0) = \lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h^2 - 0^2}{h} = \lim_{h \rightarrow 0} h = 0$   
 d  $f'(50) = \lim_{h \rightarrow 0} \frac{f(50+h) - f(50)}{h} = \lim_{h \rightarrow 0} \frac{(50+h)^2 - 50^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{100h + h^2}{h} = \lim_{h \rightarrow 0} (100 + h) = 100$   
 2 a  $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{(x+h)^2 - x^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{2xh + h^2}{h} = \lim_{h \rightarrow 0} (2x + h)$   
 b As  $h \rightarrow 0$ ,  $f'(x) = \lim_{h \rightarrow 0} (2x + h) = 2x$

- 3 a  $g = \lim_{h \rightarrow 0} \frac{(-2+h)^3 - (-2)^3}{h}$   
 $= \lim_{h \rightarrow 0} \frac{-8 + 3(-2)^2h + 3(-2)h^2 + h^3 + 8}{h}$   
 $= \lim_{h \rightarrow 0} \frac{12h - 6h^2 + h^3}{h} = \lim_{h \rightarrow 0} (12 - 6h + h^2)$   
 b  $g = 12$   
 4 a Gradient of AB =  $\frac{(-1+h)^3 - 5(-1+h) - 4}{(-1+h) - (-1)}$   
 $= \frac{-1 + 3h = 3h^2 + h^3 + 5 - 5h - 4}{h}$   
 $= \frac{h^3 - 3h^2 - 2h}{h} = h^2 - 3h - 2$   
 b gradient =  $-2$   
 5  $\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{6(x+h) - 6x}{h} = \lim_{h \rightarrow 0} \frac{6h}{h} = 6$   
 6  $\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{4(x+h)^2 - 4x^2}{h} = \lim_{h \rightarrow 0} \frac{8xh + 4h^2}{h}$   
 $= \lim_{h \rightarrow 0} (8x + 4h) = 8x$   
 7  $\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{a(x+h)^2 - ax^2}{h} = \lim_{h \rightarrow 0} \frac{(a-a)x^2 + 2axh + ah^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{2axh + ah^2}{h} = \lim_{h \rightarrow 0} (2ax + ah) = 2ax$

**Challenge**

- a  $f'(x) = \lim_{h \rightarrow 0} \frac{\frac{1}{x+h} - \frac{1}{x}}{h} = \lim_{h \rightarrow 0} \frac{x - (x+h)}{xh(x+h)} = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)}$   
 $= \lim_{h \rightarrow 0} \frac{-1}{x^2 + xh}$   
 b  $f'(x) = \lim_{h \rightarrow 0} \frac{-1}{x(x+h)} = \frac{-1}{x^2 + xh} = \frac{-1}{x^2 + 0} = \frac{-1}{x^2}$


**Exercise 12C**

- 1 a  $7x^6$       b  $8x^7$       c  $4x^3$       d  $\frac{1}{3}x^{-\frac{2}{3}}$   
 e  $\frac{1}{4}x^{-\frac{3}{4}}$       f  $\frac{1}{3}x^{-\frac{2}{3}}$       g  $-3x^{-4}$       h  $-4x^{-5}$   
 i  $-2x^{-3}$       j  $-5x^{-6}$       k  $-\frac{1}{2}x^{-\frac{3}{2}}$       l  $-\frac{1}{3}x^{-\frac{1}{3}}$   
 m  $9x^8$       n  $5x^4$       o  $3x^2$       p  $-2x^{-3}$   
 q 1      r  $3x^2$   
 2 a  $6x$       b  $54x^8$       c  $2x^3$       d  $5x^{-\frac{1}{4}}$   
 e  $\frac{15}{2}x^{\frac{1}{4}}$       f  $-10x^{-2}$       g  $6x^2$       h  $\frac{1}{2x^5}$   
 i  $x^{-\frac{1}{2}}$       j  $\frac{15}{2}\sqrt{x}$   
 3 a  $\frac{3}{4}$       b  $\frac{1}{2}$       c 3      d 2  
 4  $\frac{dy}{dx} = \frac{3}{2\sqrt{x}}$

**Exercise 12D**

- 1 a  $4x - 6$       b  $x + 12$       c  $8x$       d  $16x + 7$   
 e  $4 - 10x$   
 2 a 12      b 6      c 7      d  $2\frac{1}{2}$   
 e  $-2$       f 4  
 3 4, 0  
 4  $(-1, -8)$   
 5 1,  $-1$   
 6 6,  $-4$





c At the turning point, the gradient of  $y = f(x)$  is zero, i.e.  $f'(x) = 0$ .

**Exercise 12E**

- 1 a  $4x^3 - x^{-2}$     b  $10x^4 - 6x^{-3}$     c  $9x^{\frac{1}{2}} - x^{-\frac{3}{2}}$   
 2 a 0    b  $11\frac{1}{2}$   
 3 a  $(2\frac{1}{2}, -6\frac{1}{4})$     b  $(4, -4)$  and  $(2, 0)$   
 c  $(16, -31)$     d  $(\frac{1}{2}, 4)$   $(-\frac{1}{2}, -4)$   
 4 a  $x^{\frac{1}{2}}$     b  $-6x^{-3}$     c  $-x^{-4}$   
 d  $\frac{4}{3}x^3 - 2x^2$     e  $\frac{1}{2}x^{\frac{1}{2}} - 6x^{-4}$     f  $\frac{1}{3}x^{-\frac{2}{3}} - \frac{1}{2}x^{-2}$   
 g  $-3x^{-2}$     h  $3 + 6x^{-2}$     i  $5x^{\frac{3}{2}} + \frac{3}{2}x^{-\frac{1}{2}}$   
 j  $3x^2 - 2x + 2$     k  $12x^3 + 18x^2$     l  $24x - 8 + 2x^{-2}$   
 5 a 1    b  $\frac{2}{9}$     c -4    d 4  
 6  $-\frac{3}{4}\sqrt{2}$   
 7 a  $512 - 2304x + 4608x^2$   
 b  $f'(x) \approx \frac{d}{dx}(512 - 2304x + 4608x^2)$   
 $= -2304 + 2 \times 4608x$   
 $= 9216x - 2304$

**Exercise 12F**

- 1 a  $y + 3x - 6 = 0$     b  $4y - 3x - 4 = 0$   
 c  $3y - 2x - 18 = 0$     d  $y = x$   
 e  $y = 12x + 14$     f  $y = 16x - 22$   
 2 a  $7y + x - 48 = 0$     b  $17y + 2x - 212 = 0$   
 3  $(1\frac{2}{9}, 1\frac{8}{9})$   
 4  $y = -x, 4y + x - 9 = 0; (-3, 3)$   
 5  $y = -8x + 10, 8y - x - 145 = 0$   
 6  $(-\frac{3}{4}, \frac{9}{8})$

**Challenge**

$L$  has equation  $y = 12x - 8$ .

**Exercise 12G**




- 1 a  $x \geq -\frac{4}{3}$     b  $x \leq \frac{2}{3}$     c  $x \leq -2$   
 d  $x \leq 2, x \geq 3$     e  $x \in \mathbb{R}$     f  $x \in \mathbb{R}$   
 g  $x \geq 0$     h  $x \geq 6$   
 2 a  $x \leq 4.5$     b  $x \geq 2.5$     c  $x \geq -1$   
 d  $-1 \leq x \leq 2$     e  $-3 \leq x \leq 3$     f  $-5 \leq x \leq 5$   
 g  $0 < x \leq 9$     h  $-2 \leq x \leq 0$   
 3  $f'(x) = -6x^2 - 3$   
 $x^2 \geq 0$  for all  $x \in \mathbb{R}$ , so  $-6x^2 - 3 \leq 0$  for all  $x \in \mathbb{R}$ .  
 $\therefore f$  is decreasing for all  $x \in \mathbb{R}$ .  
 4 a Any  $p \geq 2$   
 b No. Can be any  $p \geq 2$ .



**Exercise 12H**

- 1 a  $24x + 3, 24$   
 b  $15 - 3x^2, 6x^{-3}$   
 c  $\frac{9}{2}x^{-\frac{1}{2}} + 6x^{-3}, -\frac{9}{4}x^{-\frac{3}{2}} - 18x^{-4}$   
 d  $30x + 2, 30$   
 e  $-3x^{-2} - 16x^{-3}, 6x^{-3} + 48x^{-4}$   
 2 Acceleration  $= \frac{3}{4}t^{-\frac{1}{2}} + \frac{3}{2}t^{-\frac{3}{2}}$   
 3  $\frac{3}{2}$   
 4  $-\frac{1}{2}$


**Exercise 12I**

- 1 a -28    b -17    c  $-\frac{1}{5}$   
 2 a 10    b 4    c 12.25  
 3 a  $(-\frac{3}{4}, -\frac{9}{4})$  minimum  
 b  $(\frac{1}{2}, 9\frac{1}{4})$  maximum  
 c  $(-\frac{1}{3}, 1\frac{5}{27})$  maximum,  $(1, 0)$  minimum  
 d  $(3, -18)$  minimum,  $(-\frac{1}{3}, \frac{14}{27})$  maximum  
 e  $(1, 2)$  minimum,  $(-1, -2)$  maximum  
 f  $(3, 27)$  minimum  
 g  $(\frac{9}{4}, -\frac{9}{4})$  minimum  
 h  $(2, -4\sqrt{2})$  minimum  
 i  $(\sqrt{6}, -36)$  minimum,  $(-\sqrt{6}, -36)$  minimum,  $(0, 0)$  maximum  
 4 a









5 (1, 1) inflection (gradient is positive either side of point)







6 Maximum value is 27;  $f(x) \leq 27$

7 a (1, -3): minimum, (-3, -35): minimum,  $(-\frac{1}{4}, \frac{357}{256})$ : maximum



**Exercise 12J**



c  $f(x) = x^3 - 7x^2 + 8x + 16$

$f'(x) = 3x^2 - 14x + 8 = (3x - 2)(x - 4)$

d  $(4, 0)$ ,  $(\frac{2}{3}, 0)$  and  $(0, 8)$





**Exercise 12K**

1  $2t - 3$       2  $2\pi$       3  $\frac{4}{3}$

4  $48\pi$       5  $18$

6 a Let  $x$  = width of garden.

$x + 2y = 80$

$A = xy = 2y$

b  $20\text{ m} \times 40\text{ m}, 800\text{ m}^2$

7 a  $2\pi r^2 + 2\pi r h = 600\pi \Rightarrow h = \frac{300 - r^2}{r}$

$V = \pi r^2 h = \pi r(300 - r^2) = 300\pi r - \pi r^3$

b  $2000\pi\text{ cm}^3$

8 a Let  $\theta$  = angle of sector.

$\pi r^2 \times \frac{\theta}{360} = 100 \Rightarrow \theta = \frac{36000}{\pi r^2}$

$P = 2r + 2\pi r \times \frac{\theta}{360} = 2r + \frac{200\pi r}{\pi r^2}$

$= 2r + \frac{100}{r}$

$\theta < 2\pi \Rightarrow \text{Area} < \pi r^2$ , so  $\pi r^2 > 100$

$\therefore r > \sqrt{\frac{100}{\pi}}$

b  $40\text{ cm}$

9 a Let  $h$  = height of rectangle.

$P = \pi r + 2r + 2h = 40 \Rightarrow 2h = 40 - 2r - \pi r$

$A = \frac{\pi}{2}r^2 + 2rh = \frac{\pi}{2}r^2 + r(40 - 2r - \pi r)$

$= 40r - 2r^2 - \frac{\pi}{2}r^2$

b  $\frac{800}{4 + \pi}\text{ cm}^2$

10 a  $18x + 14y = 1512 \Rightarrow y = \frac{1512 - 18x}{14}$

$A = 12xy = 12x\left(\frac{1512 - 18x}{14}\right)$

$= 1296x - \frac{108x^2}{7}$

b  $27\,216\text{ mm}^2$

**Mixed exercise**

1  $f'(x) = \lim_{h \rightarrow 0} \frac{10(x+h)^2 - 10x^2}{h} = \lim_{h \rightarrow 0} \frac{20xh + 10h^2}{h}$   
 $= \lim_{h \rightarrow 0} (20x + 10h) = 20x$

2 a  $y$ -coordinate of  $B = (\delta x)^3 + 3(\delta x)^2 + 6\delta x + 4$

Gradient  $= \frac{((\delta x)^3 + 3(\delta x)^2 + 6\delta x + 4) - 4}{(1 + \delta x) - 1}$

$= \frac{(\delta x)^3 + 3(\delta x)^2 + 6\delta x}{(\delta x)} = (\delta x)^2 + 3\delta x + 6$

b  $6$

3  $4, 11\frac{3}{4}, 17\frac{25}{27}$

4  $2, 2\frac{2}{3}$

5  $(2, -13)$  and  $(-2, 15)$

6 a  $1 - \frac{9}{x^2}$       b  $x = \pm 3$

7  $\frac{3}{2}x^{-\frac{1}{2}} + 2x^{-\frac{3}{2}}$

8 a  $\frac{dy}{dx} = 6x^{-\frac{1}{2}} - \frac{3}{2}x^{-\frac{3}{2}} = \frac{3}{2}x^{-\frac{1}{2}}(4 - x)$       b  $(4, 16)$

9 a  $x + x^{\frac{3}{2}} - x^{-\frac{1}{2}} - 1$       b  $1 + \frac{3}{2}x^{\frac{1}{2}} + \frac{1}{2}x^{-\frac{3}{2}}$       c  $4\frac{1}{16}$

10  $6x^2 + \frac{1}{2}x^{-\frac{1}{2}} - 2x^{-2}$

11  $a = 1, b = -4, c = 5$

12 a  $3x^2 - 10x + 5$

b i  $\frac{1}{3}$       ii  $y = 2x - 7$       iii  $\frac{7}{2}\sqrt{5}$

13  $y = 9x - 4$  and  $9y + x = 128$

14 a  $(\frac{4}{5}, -\frac{2}{5})$       b  $\frac{1}{5}$

15  $P$  is  $(0, -1)$ ,  $\frac{dy}{dx} = 3x^2 - 4x - 4$

Gradient at  $P = -4$ , so  $L$  is  $y = -4x - 1$ .

$-4x - 1 = x^3 - 2x^2 - 4x - 1 \Rightarrow x^2(x - 2) = 0$

$x = 2 \Rightarrow y = -9$ , so  $Q$  is  $(2, -9)$

Distance  $PQ = \sqrt{(2 - 0)^2 + (-9 - (-1))^2} = \sqrt{68} = 2\sqrt{17}$

16 a  $x = 4, y = 20$

b  $\frac{d^2y}{dx^2} = \frac{3}{4}x^{-\frac{1}{2}} + 96x^3$

At  $x = 4$ ,  $\frac{d^2y}{dx^2} = \frac{15}{8} > 0$

$(4, 20)$  is a local minimum.


17  $(1, -11)$  and  $(\frac{7}{3}, -\frac{329}{27})$

18 a  $7\frac{31}{32}$

b  $f'(x) = (x - \frac{1}{x})^2 \geq 0$  for all values of  $x$

19  $(1, 4)$

20 a  $(1, 33)$  maximum,  $(2, 28)$  and  $(-1, 1)$  minimum



21 a  $\frac{250}{x^2} - 2x$       b  $(5, 125)$


22 a  $P(x, 5 - \frac{1}{2}x^2)$   
 $OP^2 = (x - 0)^2 + (5 - \frac{1}{2}x^2 - 0)^2$   
 $= \frac{1}{4}x^4 - 4x^2 + 25$

b  $x = \pm 2\sqrt{2}$  or  $x = 0$

c  $OP = 3$ ;  $f'(x) \geq 0$  so minimum when  $x = \pm 2\sqrt{2}$ , maximum when  $x = 0$

23 a  $3 + 5(3) + 3^2 - 3^3 = 0$  therefore  $C$  on curve

b  $A$  is  $(-1, 0)$ ;  $B$  is  $(\frac{5}{3}, 9\frac{13}{27})$



25  $\frac{10}{3}, \frac{2300\pi}{27}$

26  $\frac{dA}{dx} = 4\pi x - \frac{2000}{x^2}$

$\frac{dA}{dx} = 0: 4\pi x = \frac{2000}{x^2} \rightarrow x^3 = \frac{2000}{4\pi} = \frac{500}{\pi}$

27 a  $y = 1 - \frac{x}{2} - \frac{\pi x}{4}$

$$\begin{aligned} \text{b } R &= xy + \frac{\pi}{2} \left(\frac{x}{2}\right)^2 \\ &= x \left(1 - \frac{x}{2} - \frac{\pi x}{4}\right) + \frac{\pi x^2}{8} \\ &= x - \frac{x^2}{2} - \frac{\pi x^2}{4} + \frac{\pi x^2}{8} \\ &= \frac{x}{8}(8 - 4x - \pi x) \end{aligned}$$

$$\text{c } \frac{2}{4 + \pi} \text{ m}^2 \text{ (0.280 m}^2\text{)}$$

$$28 \text{ a } \pi x^2 + 2\pi x + \pi x^2 + 2\pi x h = 80\pi$$

$$h = \frac{40 - x - x^2}{x}$$

$$\begin{aligned} V &= \pi x^2 h = \pi x^2 \left(\frac{40 - x - x^2}{x}\right) \\ &= \pi(40x - x^2 - x^3) \end{aligned}$$

$$\text{b } \frac{10}{3} \quad \text{c } \frac{d^2V}{dx^2} < 0 \therefore \text{maximum}$$

$$\text{d } \frac{2300\pi}{27} \quad \text{e } 22\frac{2}{9}\%$$

$$29 \text{ a } \text{Length of short sides} = \frac{x}{\sqrt{2}}$$

$$\text{Area} = \frac{1}{2} \times \text{base} \times \text{height}$$

$$= \frac{1}{2} \left(\frac{x^2}{2}\right) = \frac{1}{4}x^2 \text{ m}^2$$

b Let  $l$  be length of  $EF$ .

$$\frac{1}{4}x^2 l = 4000 \Rightarrow l = \frac{16000}{x^2}$$

$$S = 2\left(\frac{1}{4}x^2\right) + \frac{2xl}{\sqrt{2}}$$

$$= \frac{1}{2}x^2 + \frac{32000x}{\sqrt{2}x^2} = \frac{x^2}{2} + \frac{16000\sqrt{2}}{x}$$

$$\text{c } x = 20\sqrt{2}, S = 1200 \text{ m}^2 \quad \text{d } \frac{d^2S}{dx^2} > 0$$

**Challenge**

$$\text{a } x^7 + 7x^6h + 21x^5h^2 + 35x^4h^3$$

$$\begin{aligned} \text{b } \frac{d}{dx}(x^7) &= \lim_{h \rightarrow 0} \frac{(x+h)^7 - x^7}{h} = \lim_{h \rightarrow 0} \frac{7x^6h + 21x^5h^2 + 35x^4h^3}{h} \\ &= \lim_{h \rightarrow 0} (7x^6 + 21x^5h + 35x^4h^2) = 7x^6 \end{aligned}$$


**CHAPTER 13**

**Prior knowledge check**


$$1 \text{ a } 5x^{\frac{3}{2}} \quad \text{b } 2x^{\frac{1}{2}} \quad \text{c } x^{\frac{3}{2}} - \sqrt{x} \quad \text{d } x^{-\frac{1}{2}} + 4x$$

$$2 \text{ a } 6x^2 + 3 \quad \text{b } x - 1 \quad \text{c } 3x^2 + 2x \quad \text{d } \frac{1}{x^2} - 3x^2$$

3 a



b



**Exercise 13A**

- |  |  |
|--|--|
| 1 a $y = \frac{1}{6}x^6 + c$           | b $y = 2x^5 + c$                       |
| c $y = x^{-1} + c$                     | d $y = 2x^{-2} + c$                    |
| e $y = \frac{3}{5}x^{\frac{5}{3}} + c$ | f $y = \frac{8}{3}x^{\frac{3}{2}} + c$ |
| g $y = -\frac{2}{7}x^7 + c$            | h $y = 2x^{\frac{1}{2}} + c$           |
| i $y = -10x^{-\frac{1}{2}} + c$        | j $y = \frac{9}{2}x^{\frac{3}{4}} + c$ |
| k $y = 3x^{12} + c$                    | l $y = 2x^{-7} + c$                    |
| m $y = -9x^{\frac{1}{3}} + c$          | n $y = -5x + c$                        |
| o $y = 3x^2 + c$                       | p $y = \frac{10}{3}x^{0.6} + c$        |

- |   |   |
|---|---|
| 2 a $y = \frac{1}{4}x^4 - 3x^{\frac{1}{2}} + 6x^{-1} + c$ | b $y = x^4 + 3x^{\frac{1}{3}} + x^{-1} + c$             |
| c $y = 4x + 4x^{-3} + 4x^{\frac{1}{2}} + c$               | d $y = 3x^{\frac{3}{5}} - 2x^5 - \frac{1}{2}x^{-2} + c$ |
| e $y = 4x^{-\frac{1}{3}} - 3x + 4x^2 + c$                 | f $y = x^5 + 2x^{\frac{1}{2}} + 3x^{-4} + c$            |
- 3 a  $f(x) = 6x^2 - 3x^{\frac{1}{2}} + 5x + c$     b  $f(x) = x^6 - x^{-6} + x^{\frac{1}{3}} + c$   
 c  $f(x) = x^{\frac{1}{2}} + x^{-\frac{1}{2}} + c$     d  $f(x) = 2x^5 - 4x^{-2} + c$   
 e  $f(x) = 3x^{\frac{2}{3}} - 6x^{-\frac{2}{3}} + c$   
 f  $f(x) = 3x^3 - 2x^{-2} + \frac{1}{2}x^{\frac{1}{2}} + c$

$$4 \quad y = \frac{4x^3}{3} + 6x^2 + 9x + c$$

$$5 \quad f(x) = -3x^{-1} + 4x^{\frac{3}{2}} + \frac{x^2}{2} - 4x + c$$

**Challenge**

$$y = -\frac{12}{7x^{\frac{7}{2}}} - \frac{4}{5x^{\frac{5}{2}}} + \frac{3}{2x^2} + \frac{1}{x} + c$$

**Exercise 13B**

- |                         |                        |
|-------------------------|------------------------|
| 1 a $\frac{x^4}{4} + c$ | b $\frac{x^8}{8} + c$  |
| c $-x^{-3} + c$         | d $\frac{5x^3}{3} + c$ |
- 2 a  $\frac{1}{5}x^5 + \frac{1}{2}x^4 + c$     b  $\frac{x^4}{2} - \frac{x^3}{3} + \frac{5x^2}{2} + c$   
 c  $2x^{\frac{3}{2}} - x^3 + c$
- 3 a  $-4x^{-1} + 6x^{\frac{1}{2}} + c$     b  $-6x^{-1} - \frac{2}{5}x^{\frac{5}{2}} + c$   
 c  $-4x^{-\frac{1}{2}} + \frac{x^3}{3} - 2x^{\frac{1}{2}} + c$
- 4 a  $x^4 + x^{-3} + rx + c$     b  $\frac{1}{2}x^2 + 2x^{\frac{1}{2}} - 2x^{-\frac{1}{2}} + c$   
 c  $\frac{px^5}{5} + 2tx - 3x^{-1} + c$
- 5 a  $t^3 + t^{-1} + c$     b  $\frac{2}{3}t^3 + 6t^{-\frac{1}{2}} + t + c$   
 c  $\frac{p}{4}t^4 + q^2t + px^3t + c$
- 6 a  $2x - \frac{3}{x} + c$     b  $\frac{4}{3}x^3 + 6x^2 + 9x + c$   
 c  $\frac{4}{5}x^{\frac{5}{2}} + 2x^{\frac{3}{2}} + c$



- 7 a  $\frac{1}{3}x^3 + 2x - \frac{1}{x} + c$       b  $\frac{1}{2}x^2 + \frac{8}{3}x^{\frac{3}{2}} + 4x + c$   
 c  $2x^{\frac{1}{2}} + \frac{4}{3}x^{\frac{3}{2}} + c$
- 8 a  $\frac{3}{5}x^{\frac{3}{2}} - \frac{2}{x^2} + c$       b  $-\frac{1}{x^2} - \frac{1}{x} + 3x + c$   
 c  $\frac{1}{x}x^4 - \frac{1}{3}x^3 + \frac{3}{2}x^2 - 3x + c$       d  $\frac{8}{5}x^{\frac{5}{2}} + \frac{8}{3}x^{\frac{3}{2}} + 2x^{\frac{1}{2}} + c$   
 e  $3x + 2x^{\frac{1}{2}} + 2x^{\frac{3}{2}} + c$       f  $\frac{2}{5}x^{\frac{5}{2}} + 3x^2 + 6x^{\frac{3}{2}} + c$
- 9 a  $-\frac{A}{x} - 3x + c$       b  $\frac{2}{3}\sqrt{p}x^{\frac{3}{2}} - \frac{1}{x^2} + c$   
 c  $-\frac{p}{x} + \frac{2qx^{\frac{3}{2}}}{3} + rx + c$
- 10  $-\frac{6}{x} + \frac{8x^{\frac{3}{2}}}{3} - \frac{3x^2}{2} + 2x + c$
- 11  $2x^4 + 3x^2 - 6x^{\frac{1}{2}} + c$
- 12 a  $(2 + 5\sqrt{x})^2 = 4 + 10\sqrt{x} + 10\sqrt{x} + 25x = 4 + 20\sqrt{x} + 25x$   
 b  $4x + \frac{40x^{\frac{3}{2}}}{3} + \frac{25x^2}{2} + c$
- 13  $\frac{x^6}{2} - 8x^{\frac{1}{2}} + c$
- 14  $p = -4, q = -2.5$
- 15 a  $1024 - 5120x + 11520x^2$   
 b  $1024x - 2560x^2 + 3840x^3 + c$

**Exercise 13C**

- 1 a  $y = x^3 + x^2 - 2$       b  $y = x^4 - \frac{1}{x^2} + 3x + 1$   
 c  $y = \frac{2}{3}x^{\frac{3}{2}} + \frac{1}{12}x^3 + \frac{1}{3}$       d  $y = 6\sqrt{x} - \frac{1}{2}x^2 - 4$   
 e  $y = \frac{1}{3}x^3 + 2x^2 + 4x + \frac{2}{3}$       f  $y = \frac{2}{5}x^{\frac{5}{2}} + 6x^{\frac{1}{2}} + 1$
- 2  $f(x) = \frac{1}{2}x^4 + \frac{1}{x} + \frac{1}{2}$
- 3  $y = 1 - \frac{2}{\sqrt{x}} - \frac{3}{x}$
- 4  $f(x) = 3x^3 + 2x^2 - 3x - 2$
- 5  $y = 6x^{\frac{1}{2}} - \frac{4x^{\frac{3}{2}}}{5} + \frac{118}{5}$
- 6 a  $p = \frac{1}{2}, q = 1$       b  $y = 4x^{\frac{3}{2}} + \frac{5x^2}{2} - \frac{421}{2}$
- 7 a  $f(t) = 10t - \frac{5t^2}{2}$       b  $7\frac{1}{2}$
- 8 a  $f(t) = -4.9t^2 + 35$       b 23.975 m  
 c 35 m      d 2.67 seconds  
 e e.g. the ground is flat

**Challenge**

- 1  $f_2(x) = \frac{x^3}{3}; f_4(x) = \frac{x^4}{12}$       b  $\frac{x^{n+1}}{3 \times 4 \times 5 \times \dots \times (n+1)}$
- 2  $f_2(x) = x + 1; f_3(x) = \frac{1}{2}x^2 + x + 1; f_4(x) = \frac{1}{6}x^3 + \frac{1}{2}x^2 + x + 1$

**Exercise 13D**

- 1 a  $152\frac{1}{4}$       b  $48\frac{2}{5}$       c  $5\frac{1}{3}$       d 2  
 2 a  $5\frac{1}{4}$       b 10      c  $11\frac{5}{6}$       d  $60\frac{1}{2}$   
 3 a  $16\frac{2}{3}$       b  $46\frac{1}{2}$       c  $\frac{11}{14}$       d  $2\frac{1}{2}$   
 4  $A = -7$  or 4  
 5 28  
 6  $-8 + 8\sqrt{3}$   
 7  $k = \frac{25}{4}$   
 8 450 m

**Challenge**



$k = 2$

**Exercise 13E**

- 1 a 22      b  $36\frac{2}{3}$       c  $48\frac{8}{15}$       d 6  
 2 4      3 6      4  $10\frac{2}{3}$   
 5  $21\frac{1}{3}$       6  $\frac{4}{81}$       7  $k = 2$   
 8 a  $(-1, 0)$  and  $(3, 0)$       b  $10\frac{2}{3}$   
 9  $1\frac{1}{3}$



**Exercise 13F**

- 1 a  $1\frac{1}{3}$       b  $20\frac{5}{6}$




- c  $40\frac{1}{2}$

- d  $1\frac{1}{3}$



- e  $21\frac{1}{12}$



- 2 a  $(-3, 0)$  and  $(2, 0)$       b  $21\frac{1}{12}$   
 3 a  $f(-3) = 0$   
 b  $f(x) = (x + 3)(-x^2 + 7x - 10)$   
 c  $f(x) = (x + 3)(x - 5)(2 - x)$   
 d  $(-3, 0), (2, 0)$  and  $(5, 0)$   
 e  $143\frac{5}{6}$

**Challenge**

- 1 a  $4\frac{1}{2}$       b 9      c  $\frac{9a}{2}$       d  $4\frac{1}{2}$       e  $\frac{9}{2a}$   
 2 a B has x-coordinate 1

$$\int_0^1 (x^3 + x^2 - 2x) dx = \left[ \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2 \right]_0^1$$

$$= \frac{1}{4} + \frac{1}{3} - 1 = -\frac{5}{12}$$

So area under x-axis is  $\frac{5}{12}$

Area above x-axis is

$$\left( \frac{1}{4}0^4 + \frac{1}{3}0^3 - 0^2 \right) - \left( \frac{1}{4}x^4 + \frac{1}{3}x^3 - x^2 \right) = \frac{5}{12}$$

So the x-coordinate of a satisfies

$$3x^4 + 4x^3 - 12x^2 + 5 = 0$$

Then use the factor theorem twice to get

$$(x - 1)^2(3x^2 + 10x + 5) = 0$$

b A has coordinates  $\left(\frac{-5 + \sqrt{10}}{3}, \frac{-80 + 37\sqrt{10}}{27}\right)$

The roots at 1 correspond to point B.

The root  $\frac{-5 - \sqrt{10}}{3}$  gives a point on the curve to the left of -2 below the x-axis, so cannot be A.

**Exercise 13G**

- 1 a  $A(-2, 6), B(2, 6)$       b  $10\frac{2}{3}$
- 2 a  $A(1, 3), B(3, 3)$       b  $1\frac{1}{3}$
- 3  $6\frac{2}{3}$
- 4 4.5
- 5 a (2, 12)      b  $13\frac{1}{3}$
- 6 a  $20\frac{5}{6}$       b  $17\frac{1}{6}$
- 7 a, b Substitute into equation for y
- c  $y = x - 4$       d  $8\frac{3}{5}$
- 8  $3\frac{3}{8}$
- 9 a Substitute  $x = 4$  into both equations
- b 7.2
- 10 a  $21\frac{1}{3}$       b  $2\frac{5}{9}$
- 11 a (-1, 11) and (3, 7)      b  $21\frac{1}{3}$

**Mixed exercise**

- 1 a  $\frac{2}{9}x^3 - \frac{3}{2}x^2 - 5x + c$       b  $\frac{3}{4}x^{\frac{4}{3}} + \frac{3}{2}x^{\frac{2}{3}} + c$
- 2  $\frac{1}{3}x^3 - \frac{3}{2}x^2 + \frac{2}{x} + \frac{1}{6}$
- 3 a  $2x^4 - 2x^3 + 5x + c$       b  $2x^{\frac{5}{2}} + \frac{4}{3}x^{\frac{3}{2}} + c$
- 4  $\frac{4}{5}x^{\frac{5}{2}} - \frac{2}{3}x^{\frac{3}{2}} - 6x^{\frac{1}{2}} + c$
- 5  $x = \frac{1}{3}t^3 + t^2 + t - 8\frac{2}{3}, x = 12\frac{1}{3}$
- 6 a  $A = 6, B = 9$       b  $\frac{3}{5}x^{\frac{5}{3}} + \frac{9}{2}x^{\frac{4}{3}} + 9x + c$
- 7 a  $\frac{9}{2}x^{-\frac{1}{2}} - 8x^{-\frac{3}{2}}$       b  $6x^{\frac{2}{3}} + 32x^{\frac{1}{3}} - 24x + c$
- 8  $a = 4, b = -3.5$
- 9 25.9 m
- 10 a  $f(t) = 5t + t^2$       b 7.8 seconds
- 11 a -1, 3      b  $10\frac{2}{3}$
- 12 a  $-\frac{2x^{\frac{3}{2}}}{3} + 5x - 8\sqrt{x} + c$       b  $\frac{7}{3}$
- 13 a (3, 0)      b (1, 4)      c  $6\frac{3}{4}$
- 14 a  $\frac{3}{2}x^{-\frac{1}{2}} + 2x^{-\frac{3}{2}}$       b  $2x^{\frac{2}{3}} - 8x^{\frac{1}{3}} + c$       c  $A = 6, B = -2$
- 15 a  $\frac{dy}{dx} = 6x^{-\frac{1}{2}} - \frac{3}{2}x^{\frac{1}{2}} = \frac{3}{2}x^{-\frac{1}{2}}(4 - x)$
- b (4, 16)      c 133 (3 sf)
- 16 a (6, 12)      b  $13\frac{1}{3}$
- 17 a  $A(1, 0), B(5, 0), C(6, 5)$       b  $10\frac{1}{6}$
- 18 a  $q = -2$       b  $C(6, 17)$       c  $1\frac{1}{3}$
- 19  $-\frac{9}{x} - \frac{16x^{\frac{3}{2}}}{3} + 2x^2 - 5x + c$
- 20  $A = -6$  or 1
- 21 a  $f'(x) = \frac{(2 - x^2)(4 - 4x^2 + x^4)}{x^2} = 8x^{-2} - 12 + 6x^2 - x^4$
- b  $f''(x) = -16x^{-3} + 12x - 4x^3$
- c  $f(x) = -\frac{8}{x} - 12x + 2x^3 - \frac{x^5}{5} - \frac{47}{5}$
- 22 a  $(-3, 0)$  and  $(\frac{1}{2}, 0)$       b  $14\frac{7}{24}$
- 23 a  $(-\frac{3}{2}, 0)$  and (4, 0)      b  $55\frac{11}{24}$
- 24 a -2 and 3      b  $21\frac{1}{12}$

**Challenge**


$10\frac{5}{12}$

**CHAPTER 14**


**Prior knowledge check**

- 1 a 125      b  $\frac{1}{3}$       c 32      d 49      e 1
- 2 a  $6^6$       b  $y^{21}$       c  $2^6$       d  $x^4$
- 3 gradient 1.5, intercept 4.1


**Exercise 14A**




b  $x \approx 2.6$



b  $x \approx -1.4$




- 4 a True, because  $a^0 = 1$  whenever  $a$  is positive
- b False, for example when  $a = \frac{1}{2}$
- c True, because when  $a$  is positive,  $a^x > 0$  for all values of  $x$






- 6  $k = 3, a = 2$   
 7 a As  $x$  increases,  $y$  decreases  
 b  $p = 1.2, q = 0.2$

Challenge









Exercise 14B


- 1 a 2.7183 b 54.5982 c 0.0000 d 1.2214  
 2 a



$e = 2.71828\dots$   
 $e^3 = 20.08553\dots$



- 4 a  $A = 1, C = 5, b$  is positive  
 b  $A = 4, C = 0, b$  is negative  
 c  $A = 6, C = 2, b$  is positive  
 5  $A = e^2, b = 3$




- 6 a  $6e^{6x}$  b  $-\frac{1}{3}e^{-3x}$  c  $14e^{2x}$   
 d  $2e^{0.4x}$  e  $3e^{3x} + 2e^x$  f  $2e^{2x} + e^x$   
 7 a  $3e^6$  b 3 c  $3e^{-1.5}$   
 8  $f'(x) = 0.2e^{0.2x}$   
 The gradient of the tangent when  $x = 5$  is  
 $f'(5) = 0.2e^1 = 0.2e$ .  
 The equation of the tangent is therefore  
 $e = 0.2e \times 5 + c$ , so  $c = 0$ .

Exercise 14C


- 1 a £20 000 b £14 331  
 c
-

- 2 a 30 000      b 38 221



d Model predicts population of the country to be over 200 million, this is highly unlikely and by 2500 new factors are likely to affect population growth. Model not valid for predictions that far into the future.

- 3 a 200  
b Disease will infect up to 300 people.




- 4 a i 15 rabbits      ii 132 rabbits  
b The initial number of rabbits

c  $\frac{dR}{dm} = 2.4e^{0.2m}$   
When  $m = 6$ ,  $\frac{dR}{dm} = 7.97 \approx 8$

d The rabbits may begin to run out of food or space

- 5 a 0.565 bars  
b  $\frac{dp}{dh} = -0.13e^{-0.13h} = -0.13p$ ,  $k = -0.13$   
c The atmospheric pressure decreases exponentially as the altitude increases  
d 12%  
6 a Model 1: £15 733  
Model 2: £15 723 Similar results  
b Model 1: £1814  
Model 2: £2484 Model 2 predicts a larger value



d In Model 2 the tractor will always be worth at least £1000. This could be the value of the tractor as scrap metal.

### Exercise 14D

- 1 a  $\log_4 256 = 4$       b  $\log_3 \frac{1}{9} = -2$   
c  $\log_{10} 1\,000\,000 = 6$       d  $\log_{11} 11 = 1$   
e  $\log_{0.2} 0.008 = 3$   
2 a  $2^4 = 16$       b  $5^2 = 25$   
c  $9^{\frac{1}{2}} = 3$       d  $5^{-1} = 0.2$   
e  $10^5 = 100\,000$

- 3 a 3      b 2      c 7      d 1  
e 6      f  $\frac{1}{2}$       g -1      h -2  
i 10      j -2  
4 a 625      b 9      c 7      d 9  
e 20      f 2  
5 a 2.475      b 2.173      c 3.009      d 1.099  
6 a  $5 = \log_2 32 < \log_2 50 < \log_2 64 = 6$   
b 5.644  
7 a i 1      ii 1      iii 1      b  $a^1 \equiv a$   
8 a i 0      ii 0      iii 0      b  $a^0 \equiv 1$

### Exercise 14E


- 1 a  $\log_2 21$       b  $\log_2 9$       c  $\log_5 80$   
d  $\log_6 \left(\frac{64}{81}\right)$       e  $\log_{10} 120$   
2 a  $\log_2 8 = 3$       b  $\log_6 36 = 2$       c  $\log_{12} 144 = 2$   
d  $\log_8 2 = \frac{1}{3}$       e  $\log_{10} 10 = 1$   
3 a  $3 \log_a x + 4 \log_a y + \log_a z$   
b  $5 \log_a x - 2 \log_a y$   
c  $2 + 2 \log_a x$   
d  $\log_a x - \frac{1}{2} \log_a y - \log_a z$   
e  $\frac{1}{2} + \frac{1}{2} \log_a x$   
4 a  $\frac{4}{3}$       b  $\frac{1}{18}$       c  $\sqrt{30}$       d 2  
5 a  $\log_3(x+1) - 2 \log_3(x-1) = 1$   
 $\log_3 \left(\frac{x+1}{(x-1)^2}\right) = 1$   
 $\frac{x+1}{(x-1)^2} = 3$   
 $x+1 = 3(x-1)^2$   
 $x+1 = 3(x^2 - 2x + 1)$   
 $3x^2 - 7x + 2 = 0$   
b  $x = 2$   
6 a  $a = 9$ ,  $b = 4$

### Challenge

$\log_a x = m$  and  $\log_a y = n$   
 $x = a^m$  and  $y = a^n$   
 $x \div y = a^m \div a^n = a^{m-n}$   
 $\log_a \left(\frac{x}{y}\right) = m - n = \log_a x - \log_a y$

### Exercise 14F

- 1 a 6.23      b 2.10      c 0.431  
d 1.66      e -3.22      f 1.31  
g 1.25      h -1.73  
2 a 0, 2.32      b 1.26, 2.18      c 1.21  
d 0.631      e 0.565, 0.712      f 0  
g 2      h -1  
3 a 5.92      b 3.2  
4 a (0, 1)



- b  $\frac{1}{2}, \frac{3}{2}$   
5 a 0.7565      b 7.9248      c 0.2966





**Exercise 14G**

- 1 a  $\ln 6$       b  $\frac{1}{2} \ln 11$       c  $3 - \ln 20$   
 d  $\frac{1}{4} \ln \left(\frac{1}{3}\right)$       e  $\frac{1}{2} \ln 3 - 3$       f  $5 - \ln 19$
- 2 a  $e^2$       b  $\frac{e}{4}$       c  $\frac{1}{2} e^4 - \frac{3}{2}$   
 d  $\frac{1}{6} (e^3 + 2)$       e  $18 - e^{\frac{1}{2}}$       f 2, 5
- 3 a  $\ln 2, \ln 6$       b  $\frac{1}{2} \ln 2, 0$       c  $e^3, e^{-5}$   
 d  $\ln 4, 0$       e  $\ln 5, \ln \left(\frac{1}{3}\right)$       f  $e^6, e^{-2}$
- 4  $\ln 3, 2 \ln 2$
- 5 a  $\frac{1}{8} (e^2 + 3)$       b  $\frac{1}{5} (\ln 3 + 40)$       c  $\frac{1}{5} \ln 7, 0$   
 d  $e^3, e^{-1}$
- 6  $\frac{1 + \ln 5}{4 + \ln 3}$
- 7 a The initial concentration of the drug in mg/l  
 b 4.91 mg/l  
 c  $3 = 6e^{-\frac{t}{10}}$   
 $\frac{1}{2} = e^{-\frac{t}{10}}$   
 $\ln \left(\frac{1}{2}\right) = -\frac{t}{10}$   
 $t = -10 \ln \left(\frac{1}{2}\right) = 6.931\dots = 6 \text{ hours } 56 \text{ minutes}$
- 8 a  $(0, 3 + \ln 4)$       b  $(4 - e^{-3})$


**Challenge**

As  $y = 2$  is an asymptote,  $C = 2$ .  
 Substituting  $(0, 5)$  gives  $5 = Ae^0 + 2$ , so  $A$  is 3.  
 Substituting  $(6, 10)$  gives  $10 = 3e^{6b} + 2$ .  
 Rearranging this gives  $B = \frac{1}{6} \ln \left(\frac{8}{3}\right)$ .


**Exercise 14H**

- 1 a  $\log S = \log(4 \times 7^x)$   
 $\log S = \log 4 + \log 7^x$   
 $\log S = \log 4 + x \log 7$   
 b gradient  $\log 7$ , intercept  $\log 4$
- 2 a  $\log A = \log(6x^4)$   
 $\log A = \log 6 + \log x^4$   
 $\log A = \log 6 + 4 \log x$   
 b gradient 4, intercept  $\log 6$
- 3 a Missing values 1.52, 1.81, 1.94
- b 
- c  $a = 3.5, n = 1.4$
- 4 a Missing values 2.63, 3.61, 4.49, 5.82
- b 
- c  $b = 3.4, a = 10$

- 5 a Missing values  $-0.39, 0.62, 1.54, 2.81$




- c  $a = 60, b = 0.75$   
 d 1,600 kcal per day (2 s.f.)
- 6 a Missing values 2.94, 1.96, 0.95



- c  $A = 5800, b = -0.9$   
 d 694 times
- 7 a Missing values 0.98, 1.08, 1.13, 1.26, 1.37

- b  $P = ab^t$   
 $\log P = \log(ab^t)$   
 $\log P = \log a + \log b^t$   
 $\log P = \log a + t \log b$



- d  $a = 7.6, b = 1.0$   
 e The rate of growth is often proportional to the size of the population
- 8 a  $\log N = 0.095t + 1.6$   
 b  $a = 40, b = 1.2$   
 c The initial number of sick people  
 d 9500 people. After 30 days people may start to recover, or the disease may stop spreading as quickly.
- 9 a  $\log A = 2 \log w - 0.1049$   
 b  $q = 2, p = 0.7854$   
 c Circles:  $p$  is approximately one quarter  $\pi$ , and the width is twice the radius, so  $A = \frac{\pi}{4} w^2 = \frac{\pi}{4} (2r)^2 = \pi r^2$ .


**Challenge**

$y = 5.8 \times 0.9^x$




Mixed exercise


1 a



b



c




- 2 a  $2 \log_a p + \log_a q$       b  $\log_a p = 4, \log_a q = 1$   
 3 a  $\frac{1}{4}p$       b  $\frac{3}{4}p + 1$   
 4 a 2.26      b 1.27      c 7.02  
 5 a  $4^x - 2^{x+1} - 15 = 0$   
 $2^{2x} - 2 \times 2^x - 15 = 0$   
 $(2^x)^2 - 2 \times 2^x - 15 = 0$   
 $u^2 - 2u - 15 = 0$   
 b 2.32  
 6  $x = 6$   
 7 a  $-e^{-x}$       b  $11e^{11x}$       c  $30e^{5x}$   
 8 a  $\frac{e^8 + 5}{2}$       b  $\frac{\ln 5}{4}$       c  $-\frac{1}{2} \ln 14$       d  $\frac{3 + \sqrt{13}}{2}$   
 e 0      f  $\frac{e^4}{2}$   
 9 a £950      b £290      c 4.28 years      d £100  
 e   
 f A good model. The computer will always be worth something

10 a  $y = \left(\frac{2}{\ln 4}\right)x$


- b (0, 0) satisfies the equation of the line.  
 c 2.43  
 11 a We cannot go backwards in time  
 b 75°C  
 c 5 minutes  
 d The exponential term will always be positive, so the overall temperature will be greater than 20°C.  
 12 a  $S = aV^b$   
 $\log S = \log(aV^b)$   
 $\log S = \log a + \log(V^b)$   
 $\log S = \log a + b \log V$

b	$\log S$	1.26	1.70	2.05	2.35	2.50
	$\log V$	0.86	1.53	2.05	2.49	2.72



- d The gradient is approximately 1.5;  $a = 0.09$   
 13 a The model concerns decay, not growth

b



- c  $70 = 140e^{30k}$   
 $\frac{1}{2} = e^{30k}$   
 $\ln\left(\frac{1}{2}\right) = 30k$   
 $k = \frac{1}{30} \ln\left(\frac{1}{2}\right)$   
 $k = -\frac{1}{30} \ln(2)$ , so  $c = -\frac{1}{30}$   
 14 a 6.3 million views  
 b  $\frac{dV}{dx} = 0.4e^{0.4x}$   
 c  $9.42 \times 10^{16}$  new views per day  
 d This is too big, so the model is not valid after 100 days  
 15 a 4.2  
 b i  $1.12 \times 10^{25}$  dyne cm  
 ii  $3.55 \times 10^{26}$  dyne cm  
 c divide b ii by b i  
 16 a They exponentiated the two terms on RHS separately rather than combining them first.  
 b  $x = 2 \pm \sqrt{5}$





**Challenge**

- a  $y = 9^x = 3^{2x}$ ,  $\log_3(y) = 2x$   
 b  $y^2 = (9^x)^2 = 9^{2x}$ ,  $\log_9(y^2) = 2x$   
 c  $x = -\frac{1}{3}$  or  $x = -2$

**Review exercise 3**

- 1 -4.5  
 2  $\sqrt{7}$   
 3 a All equal to  $\sqrt{145}$   
 b  $(x-1)^2 + (y+3)^2 = 145$   
 4 a  $-2i - 8j$   
 b  $|\vec{AB}| = |\vec{AC}| = \sqrt{85}$   
 c  $|\vec{BC}| = \sqrt{68}$   
 $\cos \angle ABC = \frac{85 + 68 - 85}{2 \times \sqrt{85} \times \sqrt{68}} = \frac{1}{\sqrt{5}}$   
 5 12  
 6 a 5N      b 7  
 7  $m = 50\sqrt{3} + 30$ ,  $n = 50$   
 8 a  $\sqrt{(-75)^2 + 180^2} = 195 > 150 = \sqrt{90^2 + 120^2}$   
 b Boat A: 6.5 m/s; Boat B: 5 m/s; Both boats arrive at the same time – it is a tie.  
 9  $\lim_{h \rightarrow 0} \frac{5(x+h)^2 - 5x^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{5x^2 + 10xh + 5h^2 - 5x^2}{h}$   
 $= \lim_{h \rightarrow 0} \frac{10xh + 5h^2}{h}$   
 $= \lim_{h \rightarrow 0} 10x + 5h$   
 $= 10x$   
 10  $\frac{dy}{dx} = 12x^2 + x^{-\frac{1}{2}}$   
 11 a  $\frac{dy}{dx} = 4 + \frac{9}{2}x^{\frac{1}{2}} - 4x$   
 b Substitute  $x = 4$  into equation for  $C$   
 c Gradient of tangent = -3 so gradient of normal =  $\frac{1}{3}$   
 Substitute  $(4, 8)$  into  $y = \frac{1}{3}x + c$   
 Rearrange  $y = \frac{1}{3}x + \frac{20}{3}$   
 d  $PQ = 8\sqrt{10}$   
 12 a  $\frac{dy}{dx} = 8x - 5x^{-2}$ , at  $P$  this is 3  
 b  $y = 3x + 5$   
 c  $k = -\frac{5}{3}$   
 13 a  $P = 2$ ,  $Q = 9$ ,  $R = 4$   
 b  $3x^{\frac{1}{2}} + \frac{9}{2}x^{-\frac{1}{2}} - 2x^{-\frac{3}{2}}$   
 c When  $x = 1$ ,  $f'(x) = 5\frac{1}{2}$ , gradient of  $2y = 11x + 3$  is  $5\frac{1}{2}$ , so it is parallel with tangent  
 14  $f'(x) = 3x^2 - 24x + 48 = 3(x-4)^2 > 0$   
 15 a  $A(1,0)$  and  $B(2,0)$   
 b  $(\sqrt{2}, 2\sqrt{2} - 3)$   
 16 a  $V = \pi r^2 h = 128\pi$ , so  $h = \frac{128}{r^2}$   
 $S = 2\pi r h + 2\pi r^2 = \frac{256\pi}{r} + 2\pi r^2$   
 b  $96\pi \text{ cm}^2$

17 a  $\frac{dy}{dx} = 6x + 2x^{-\frac{1}{2}}$


b  $\frac{d^2y}{dx^2} = 6 - x^{-\frac{3}{2}}$

c  $x^3 + \frac{8}{3}x^{\frac{3}{2}} + c$

18 a  $2x^3 - 5x^2 - 12x$

b  $x(2x+3)(x-4)$

c



19  $6\frac{3}{4}$

20 4

21 a  $-x^4 + 3x^2 + 4 = (-x^2 + 4)(x^2 + 1)$ ;  $x^2 + 1 = 0$  has no real solutions; so solutions are  $A(-2, 0)$ ,  $B(2, 0)$

b 19.2

22  $4\frac{1}{2}$  square units

23 a  $P(-1, 4)$ ,  $Q(2, 1)$

b 4.5

24 a  $k = -1$ ,  $A(0, 2)$

b  $\ln 3$

25 a  $425^\circ\text{C}$       b 7.49 minutes      c  $1.64^\circ\text{C/minute}$

d The temperature can never go below  $25^\circ\text{C}$ , so cannot reach  $20^\circ\text{C}$ .


26 a -0.179      b  $x = 15$

27 a  $x = 1.55$       b  $x = 4$  or  $x = \frac{1}{2}$

28 a  $\log_p 2$       b 0.125

29 a  $x = 2$       b  $x = \ln 3$  or  $x = \ln 1 = 0$

30 a Missing values 0.88, 1.01, 1.14 and 1.29



c  $P = ab^t$

$\log P = \log(ab^t) = \log a + t \log b$

This is a linear relationship. The gradient is  $\log b$  and the intercept is  $\log a$ .

d  $a = 5.9$ ,  $b = 1.0$

**Challenge**

1 a 0

b 1

2 a  $f'(-3) = f'(2) = 0$ , so  $f'(x) = k(x+3)(x-2) = k(x^2 + x - 6)$ ; there are no other factors as  $f(x)$  is cubic

b  $2x^3 + 3x^2 - 36x - 5$

3 51.2

4 a  $f(0) = 0^3 - k(0) + 1 = 1$ ;  $g(0) = e^{2(0)} = 1$ ;  $P(0, 1)$

b  $\frac{1}{2}$

**Practice paper**

1 a  $\frac{1}{3}$       b  $5\sqrt{2}$

2  $y = \frac{2}{3}x + \frac{8}{3}$

3 a error 1:  $= -\frac{3}{\sqrt{x}} = -3x^{-\frac{1}{2}}$ , not  $-3x^{\frac{1}{2}}$

error 2:  $\left[\frac{x^5}{5} - 2x^{\frac{3}{2}} + 2x\right]_1^2 = \left(\frac{32}{5} - 2\sqrt{8} + 4\right) - \left(\frac{1}{5} - 2 + 2\right)$

not  $\left(\frac{1}{5} - 2 + 2\right) - \left(\frac{32}{5} - 2\sqrt{8} + 4\right)$

b 5.71 (3 s.f.)

4  $x = 30^\circ, 90^\circ, 150^\circ$

5 a  $2x^2(x + 3)$

b  $2x^2(x + 3) = 980 \Rightarrow 2x^3 + 6x^2 - 980 = 0$   
 $\Rightarrow x^3 + 3x^2 - 490 = 0$

c For  $f(x) = x^3 + 3x^2 - 490 = 0$ ,  $f(7) = 0$ , so  $x - 7$  is a factor of  $f(x)$  and  $x = 7$  is a solution.

d Equation becomes  $(x - 7)(x^2 + 10x + 70) = 0$   
 Quadratic has discriminant  $10^2 - 4 \times 1 \times 70 = -180$   
 So the quadratic has no real roots, and the equation has no more real solutions.

6  $x + 15y + 106 = 0$

7 a  $\log_{10} P = 0.01t + 2$

b 100, initial population


c 1.023

d Accept answers from 195 to 200

8  $1 + \cos^4 x - \sin^4 x \equiv 1 + (\cos^2 x + \sin^2 x)(\cos^2 x - \sin^2 x)$   
 $\equiv (1 + \sin^2 x) + \cos^2 x \equiv 2\cos^2 x$

9 Magnitude =  $\sqrt{29}$ , angle =  $112^\circ$  (3 s.f.)

10 a  $56.5^\circ$  (3 s.f.)      b £49.63



b -4, -1, 4

12  $x = 3$  or  $-\frac{5}{3}$

13 a  $1 - 15x + 90x^2$

b 0.859

c Greater: The next term will be subtracting from this, and future positive terms will be smaller.

14 a  $f(x) = \int(x^{\frac{1}{2}} - 1 - x^{-2})dx = -2x^{\frac{1}{2}} - x + x^{-1} + c$   
 $= -\frac{x^2 + 2\sqrt{x} - 1}{x} + c$

b  $c = \frac{5}{3} + \frac{2}{3}\sqrt{3}$

15 a  $5^2 + 1^2 - 4 \times 5 + 6 \times 1 = 12$ , so (5, 1) lies on C.  
 Centre = (2, -3), radius = 5

b  $y = -\frac{3}{4}x + \frac{19}{4}$

c  $12^{\frac{15}{32}}$

